

Plant Products as Antimicrobial Agents

MARJORIE MURPHY COWAN*

Department of Microbiology, Miami University, Oxford, Ohio 45056

INTRODUCTION	564
Brief History	565
MAJOR GROUPS OF ANTIMICROBIAL COMPOUNDS FROM PLANTS	565
Phenolics and Polyphenols	565
Simple phenols and phenolic acids	565
Quinones	568
Flavones, flavonoids, and flavonols	568
Tannins.....	569
Coumarins.....	571
Terpenoids and Essential Oils	571
Alkaloids.....	572
Lectins and Polypeptides	572
Mixtures	572
Other Compounds.....	573
EXPERIMENTAL APPROACHES	573
Extraction Methods	573
Efficacy	574
In vitro experiments	574
(i) Bacteria and fungi.....	574
(ii) Viruses	574
(iii) Protozoa and helminths	575
In vivo testing of phytochemicals	575
Clinical trials in humans.....	575
PLANTS WITH EFFICACY AGAINST HUMAN IMMUNODEFICIENCY VIRUS	577
COMMERCIAL AVAILABILITY AND SAFETY OF COMPOUNDS	577
CONCLUSIONS AND FUTURE DIRECTIONS	577
ACKNOWLEDGMENTS	577
REFERENCES	577

INTRODUCTION

“Eat leeks in March and wild garlic in May,
and all the year after the physicians may play.”

Traditional Welsh rhyme (230)

“An apple a day keeps the doctor away.”

Traditional American rhyme

Finding healing powers in plants is an ancient idea. People on all continents have long applied poultices and imbibed infusions of hundreds, if not thousands, of indigenous plants, dating back to prehistory. There is evidence that Neanderthals living 60,000 years ago in present-day Iraq used plants such as hollyhock (211, 224); these plants are still widely used in ethnomedicine around the world. Historically, therapeutic results have been mixed; quite often cures or symptom relief resulted. Poisonings occurred at a high rate, also. Currently, of the one-quarter to one-half of all pharmaceuticals dispensed in the United States having higher-plant origins, very few are intended for use as antimicrobials, since we have relied on bacterial and fungal sources for these activities. Since the advent of antibiotics in the 1950s, the use of plant derivatives as antimicrobials has been virtually nonexistent.

Clinical microbiologists have two reasons to be interested in the topic of antimicrobial plant extracts. First, it is very likely that these phytochemicals will find their way into the arsenal of antimicrobial drugs prescribed by physicians; several are already being tested in humans (see below). It is reported that, on average, two or three antibiotics derived from microorganisms are launched each year (43). After a downturn in that pace in recent decades, the pace is again quickening as scientists realize that the effective life span of any antibiotic is limited. Worldwide spending on finding new anti-infective agents (including vaccines) is expected to increase 60% from the spending levels in 1993 (7). New sources, especially plant sources, are also being investigated. Second, the public is becoming increasingly aware of problems with the overprescription and misuse of traditional antibiotics. In addition, many people are interested in having more autonomy over their medical care. A multitude of plant compounds (often of unreliable purity) is readily available over-the-counter from herbal suppliers and natural-food stores, and self-medication with these substances is commonplace. The use of plant extracts, as well as other alternative forms of medical treatments, is enjoying great popularity in the late 1990s. Earlier in this decade, approximately one-third of people surveyed in the United States used at least one “unconventional” therapy during the previous year (60). It was reported that in 1996, sales of botanical medicines increased 37% over 1995 (116). It is speculated that the American public may be reacting to overpre-

* Mailing address: Department of Microbiology, Miami University, Middletown Campus, 4200 East University Blvd., Middletown, OH 45042. Phone: (513) 727-3231. Fax: (513) 727-3367. E-mail: cowanmm@muohio.edu.

scription of sometimes toxic drugs, just as their predecessors of the 19th century (see below) reacted to the overuse of bleeding, purging, and calomel (248).

Brief History

It is estimated that there are 250,000 to 500,000 species of plants on Earth (25). A relatively small percentage (1 to 10%) of these are used as foods by both humans and other animal species. It is possible that even more are used for medicinal purposes (146). Hippocrates (in the late fifth century B.C.) mentioned 300 to 400 medicinal plants (195). In the first century A.D., Dioscorides wrote *De Materia Medica*, a medicinal plant catalog which became the prototype for modern pharmacopoeias. The Bible offers descriptions of approximately 30 healing plants. Indeed, frankincense and myrrh probably enjoyed their status of great worth due to their medicinal properties. Reported to have antiseptic properties, they were even employed as mouthwashes. The fall of ancient civilizations forestalled Western advances in the understanding of medicinal plants, with much of the documentation of plant pharmaceuticals being destroyed or lost (211). During the Dark Ages, the Arab world continued to excavate their own older works and to build upon them. Of course, Asian cultures were also busy compiling their own pharmacopoeia. In the West, the Renaissance years saw a revival of ancient medicine, which was built largely on plant medicinals.

North America's history of plant medicinal use follows two strands—their use by indigenous cultures (Native Americans), dating from prehistory (243), and an "alternative" movement among Americans of European origin, beginning in the 19th century. Native American use of plant medicinals has been reviewed extensively in a series of articles by Moerman (146). He reported that while 1,625 species of plants have been used by various Native American groups as food, 2,564 have found use as drugs (116). According to his calculations, this leaves approximately 18,000 species of plants which were used for neither food nor drugs. Speculations as to how and why a selected number of plant species came into use for either food or drugs is fascinating but outside the scope of this review.

Among Europeans living in the New World, the use of botanicals was a reaction against invasive or toxic mainstream medicinal practices of the day. No less a luminary than Oliver Wendell Holmes noted that medical treatments in the 1800s could be dangerous and ineffective. Examples include the use of mercury baths in London "barber shops" to treat syphilis and dangerous hallucinogens as a tuberculosis "cure." In 1861 Holmes wrote, "If the whole materia medica as now used could be sunk to the bottom of the sea, it would be all the better for mankind—and all the worse for the fishes" (92). In 1887, alternative practitioners compiled their own catalogs, notably the *Homeopathic Pharmacopoeia of the United States*.

Mainstream medicine is increasingly receptive to the use of antimicrobial and other drugs derived from plants, as traditional antibiotics (products of microorganisms or their synthesized derivatives) become ineffective and as new, particularly viral, diseases remain intractable to this type of drug. Another driving factor for the renewed interest in plant antimicrobials in the past 20 years has been the rapid rate of (plant) species extinction (131). There is a feeling among natural-products chemists and microbiologists alike that the multitude of potentially useful phytochemical structures which could be synthesized chemically is at risk of being lost irretrievably (25). There is a scientific discipline known as ethnobotany (or ethnopharmacology), whose goal is to utilize the impressive array of knowledge assembled by indigenous peoples about the plant

and animal products they have used to maintain health (77, 182, 203, 234). Lastly, the ascendancy of the human immunodeficiency virus (HIV) has spurred intensive investigation into the plant derivatives which may be effective, especially for use in underdeveloped nations with little access to expensive Western medicines. (For a comprehensive review of the search for new anti-HIV agents, see reference 51.)

This review describes the current state of plant antimicrobials in the United States, ranging from extracts commonly in use, largely by the lay community, to substances being prospectively and tested by researchers and clinicians. Also, compounds potentially effective in treating HIV infections, which are being sought far and wide for probable use in North America, are also addressed. An attempt is also made to summarize the current state of knowledge of relatively undefined herbal products, since clinicians in this country will encounter their use among patients in the United States. This review does not address the worldwide use of plants as medicinals, but, rather, it focuses on plants and their extracts currently in use in North America. Only phytochemicals reported to have anti-infective properties are examined. The many plants used as immune system boosters are outside the purview of this review. Within these boundaries, reports cited in the peer-reviewed literature are given heaviest emphasis. Detailed descriptions of individual plant-derived medications can be found in the first edition of the *Physician's Desk Reference for Herbal Medicines*, published by Medical Economics Company in 1998.

MAJOR GROUPS OF ANTIMICROBIAL COMPOUNDS FROM PLANTS

Plants have an almost limitless ability to synthesize aromatic substances, most of which are phenols or their oxygen-substituted derivatives (76). Most are secondary metabolites, of which at least 12,000 have been isolated, a number estimated to be less than 10% of the total (195). In many cases, these substances serve as plant defense mechanisms against predation by microorganisms, insects, and herbivores. Some, such as terpenoids, give plants their odors; others (quinones and tannins) are responsible for plant pigment. Many compounds are responsible for plant flavor (e.g., the terpenoid capsaicin from chili peppers), and some of the same herbs and spices used by humans to season food yield useful medicinal compounds (Table 1).

Useful antimicrobial phytochemicals can be divided into several categories, described below and summarized in Table 2.

Phenolics and Polyphenols

Simple phenols and phenolic acids. Some of the simplest bioactive phytochemicals consist of a single substituted phenolic ring. Cinnamic and caffeic acids are common representatives of a wide group of phenylpropane-derived compounds which are in the highest oxidation state (Fig. 1).

The common herbs tarragon and thyme both contain caffeic acid, which is effective against viruses (245), bacteria (31, 224), and fungi (58).

Catechol and pyrogallol both are hydroxylated phenols, shown to be toxic to microorganisms. Catechol has two -OH groups, and pyrogallol has three. The site(s) and number of hydroxyl groups on the phenol group are thought to be related to their relative toxicity to microorganisms, with evidence that increased hydroxylation results in increased toxicity (76). In addition, some authors have found that more highly oxidized phenols are more inhibitory (192, 231). The mechanisms thought to be responsible for phenolic toxicity to microorgan-

TABLE 1. Plants containing antimicrobial activity^a

Common name	Scientific name	Compound	Class	Activity ^d	Relative toxicity ^b	Reference(s) ^c
Alfalfa	<i>Medicago sativa</i>	?		Gram-positive organisms	2.3	
Allspice	<i>Pimenta dioica</i>	Eugenol	Essential oil	General	2.5	
Aloe	<i>Aloe barbadensis</i> , <i>Aloe vera</i>	Latex	Complex mixture	<i>Corynebacterium</i> , <i>Salmonella</i> , <i>Streptococcus</i> , <i>S. aureus</i>	2.7	136
Apple	<i>Malus sylvestris</i>	Phloretin	Flavonoid derivative	General	3.0	101
Ashwagandha	<i>Withania somniferum</i>	Withafarin A	Lactone	Bacteria, fungi	0.0	
Aveloz	<i>Euphorbia tirucalli</i>	?		<i>S. aureus</i>	0.0	
Bael tree	<i>Aegle marmelos</i>	Essential oil	Terpenoid	Fungi		179
Balsam pear	<i>Momordica charantia</i>	?		General	1.0	
Barberry	<i>Berberis vulgaris</i>	Berberine	Alkaloid	Bacteria, protozoa	2.0	140, 163
Basil	<i>Ocimum basilicum</i>	Essential oils	Terpenoids	<i>Salmonella</i> , bacteria	2.5	241
Bay	<i>Laurus nobilis</i>	Essential oils	Terpenoids	Bacteria, fungi	0.7	
Betel pepper	<i>Piper betel</i>	Catechols, eugenol	Essential oils	General	1.0	
Black pepper	<i>Piper nigrum</i>	Piperine	Alkaloid	Fungi, <i>Lactobacillus</i> , <i>Micrococcus</i> , <i>E. coli</i> , <i>E. faecalis</i>	1.0	78
Blueberry	<i>Vaccinium</i> spp.	Fructose	Monosaccharide	<i>E. coli</i>		158
Brazilian pepper tree	<i>Schinus terebinthifolius</i>	Terebinthone	Terpenoids	General	1.0	
Buchu	<i>Barosma setulina</i>	Essential oil	Terpenoid	General	2.0	
Burdock	<i>Arctium lappa</i>		Polyacetylene, tannins, terpenoids	Bacteria, fungi, viruses	2.3	
Buttercup	<i>Ranunculus bulbosus</i>	Protoanemonin	Lactone	General	2.0	
Caraway	<i>Carum carvi</i>		Coumarins	Bacteria, fungi, viruses		24, 26, 83, 193
Cascara sagrada	<i>Rhamnus purshiana</i>	Tannins	Polyphenols Anthraquinone	Viruses, bacteria, fungi	1.0	
Cashew	<i>Anacardium pulsatilla</i>	Salicylic acids	Polyphenols	<i>P. acnes</i> Bacteria, fungi		91
Castor bean	<i>Ricinus communis</i>	?		General	0.0	
Ceylon cinnamon	<i>Cinnamomum verum</i>	Essential oils, others	Terpenoids, tannins	General	2.0	
Chamomile	<i>Matricaria chamomilla</i>	Anthemic acid	Phenolic acid	<i>M. tuberculosis</i> , <i>S. typhi-</i> <i>murium</i> , <i>S. aureus</i> , helminths	2.3	26, 83, 193
Chapparal	<i>Larrea tridentata</i>	—	Coumarins Lignan	Viruses Skin bacteria		24
Chili peppers, paprika	<i>Capsicum annuum</i>	Capsaicin	Terpenoid	Bacteria	2.0	42, 107
Clove	<i>Syzygium aromaticum</i>	Eugenol	Terpenoid	General	1.7	
Coca	<i>Erythroxylum coca</i>	Cocaine	Alkaloid	Gram-negative and -positive cocci	0.5	
Cockle	<i>Agrostemma githago</i>	?		General	1.0	
Coltsfoot	<i>Tussilago farfara</i>	?		General	2.0	
Coriander, cilantro	<i>Coriandrum sativum</i>	?		Bacteria, fungi		
Cranberry	<i>Vaccinium</i> spp.	Fructose Other	Monosaccharide	Bacteria		17, 158, 159
Dandelion	<i>Taraxacum officinale</i>	?		<i>C. albicans</i> , <i>S. cerevisiae</i>	2.7	
Dill	<i>Anethum graveolens</i>	Essential oil	Terpenoid	Bacteria	3.0	
Echinacea	<i>Echinaceae</i> <i>angustifolia</i>	?		General		153
Eucalyptus	<i>Eucalyptus globulus</i>	Tannin —	Polyphenol Terpenoid	Bacteria, viruses	1.5	
Fava bean	<i>Vicia faba</i>	Fabatin	Thionin	Bacteria		
Gamboge	<i>Garcinia hanburyi</i>		Resin	General	0.5	
Garlic	<i>Allium sativum</i>	Allicin, ajoene	Sulfoxide Sulfated terpenoids	General		150, 187, 188 250
Ginseng	<i>Panax notoginseng</i>		Saponins	<i>E. coli</i> , <i>Sporothrix schenckii</i> , <i>Staphylococcus</i> , <i>Tricho-</i> <i>phyton</i>	2.7	
Glory lily	<i>Gloriosa superba</i>	Colchicine	Alkaloid	General	0.0	
Goldenseal	<i>Hydrastis canadensis</i>	Berberine, hydrastine	Alkaloids	Bacteria, <i>Giardia duodenale</i> , trypanosomes Plasmodia	2.0	73 163
Gotu kola	<i>Centella asiatica</i>	Asiatococside	Terpenoid	<i>M. leprae</i>	1.7	
Grapefruit peel	<i>Citrus paradisa</i>		Terpenoid	Fungi		209

Continued on following page

TABLE 1—Continued

Common name	Scientific name	Compound	Class	Activity ^d	Relative toxicity ^b	Reference(s) ^c
Green tea	<i>Camellia sinensis</i>	Catechin	Flavonoid	General	2.0	
				<i>Shigella</i>		235
				<i>Vibrio</i>		226
				<i>S. mutans</i>		166
				Viruses		113
Harmel, rue	<i>Peganum harmala</i>	?		Bacteria, fungi	1.0	
Hemp	<i>Cannabis sativa</i>	β-Resericyclic acid	Organic acid	Bacteria and viruses	1.0	
Henna	<i>Lawsonia inermis</i>	Gallic acid	Phenolic	<i>S. aureus</i>	1.5	
Hops	<i>Humulus lupulus</i>	Lupulone, humulone	Phenolic acids	General	2.3	
		—	(Hemi)terpenoids			
Horseradish	<i>Armoracia rusticana</i>		Terpenoids	General		
Hyssop	<i>Hyssopus officinalis</i>	—	Terpenoids	Viruses		
(Japanese) herb	<i>Rabdosia trichocarpa</i>	Trichorabdal A	Terpene	<i>Helicobacter pylori</i>		108
Lantana	<i>Lantana camara</i>	?		General	1.0	
—	<i>Lawsonia</i>	Lawsonone	Quinone	<i>M. tuberculosis</i>	—	
Lavender-cotton	<i>Santolina chamaecyparissus</i>	?		Gram-positive bacteria, <i>Candida</i>	1.0	213
Legume (West Africa)	<i>Millettia thonningii</i>	Alpinumisoflavone	Flavone	<i>Schistosoma</i>		175
Lemon balm	<i>Melissa officinalis</i>	Tannins	Polyphenols	Viruses		245
Lemon verbena	<i>Aloysia triphylla</i>	Essential oil	Terpenoid	<i>Ascaris</i>	1.5	
			?	<i>E. coli</i> , <i>M. tuberculosis</i> , <i>S. aureus</i>		
Licorice	<i>Glycyrrhiza glabra</i>	Glabrol	Phenolic alcohol	<i>S. aureus</i> , <i>M. tuberculosis</i>	2.0	
Lucky nut, yellow	<i>Thevetia peruviana</i>	?		<i>Plasmodium</i>	0.0	
Mace, nutmeg	<i>Myristica fragrans</i>	?		General	1.5	
Marigold	<i>Calendula officinalis</i>	?		Bacteria	2.7	
Mesquite	<i>Prosopis juliflora</i>	?		General	1.5	
Mountain tobacco	<i>Arnica montana</i>	Helanins	Lactones	General	2.0	
Oak	<i>Quercus rubra</i>	Tannins	Polyphenols			
		Quercetin (available commercially)	Flavonoid			113
Olive oil	<i>Olea europaea</i>	Hexanal	Aldehyde	General		120
Onion	<i>Allium cepa</i>	Allicin	Sulfoxide	Bacteria, <i>Candida</i>		239
Orange peel	<i>Citrus sinensis</i>	?	Terpenoid	Fungi		209
Oregon grape	<i>Mahonia aquifolia</i>	Berberine	Alkaloid	<i>Plasmodium</i>	2.0	163
				Trypanosomes, general		73
Pao d'arco	<i>Tabebuia</i>	Sesquiterpenes	Terpenoids	Fungi	1.0	
Papaya	<i>Carica papaya</i>	Latex	Mix of terpenoids, organic acids, alkaloids	General	3.0	34, 168, 191
Pasque-flower	<i>Anemone pulsatilla</i>	Anemonins	Lactone	Bacteria	0.5	
Peppermint	<i>Mentha piperita</i>	Menthol	Terpenoid	General		
Periwinkle	<i>Vinca minor</i>	Reserpine	Alkaloid	General	1.5	
Peyote	<i>Lophophora williamsii</i>	Mescaline	Alkaloid	General	1.5	
Poinsettia	<i>Euphorbia pulcherrima</i>	?		General	0.0	
Poppy	<i>Papaver somniferum</i>	Opium	Alkaloids and others	General	0.5	
Potato	<i>Solanum tuberosum</i>	?		Bacteria, fungi	2.0	
Prostrate knotweed	<i>Polygonum aviculare</i>	?		General	2.0	
Purple prairie clover	<i>Petalostemum</i>	Petalostemumol	Flavonol	Bacteria, fungi		100
Quinine	<i>Cinchona</i> sp.	Quinine	Alkaloid	<i>Plasmodium</i> spp.	2.0	
Rauwolfia, chandra	<i>Rauwolfia serpentina</i>	Reserpine	Alkaloid	General	1.0	
Rosemary	<i>Rosmarinus officinalis</i>	Essential oil	Terpenoid	General	2.3	
Sainfoin	<i>Onobrychis viciifolia</i>	Tannins	Polyphenols	Ruminal bacteria		105
Sassafras	<i>Sassafras albidum</i>	?		Helminths	2.0	
Savory	<i>Satureja montana</i>	Carvacrol	Terpenoid	General	2.0	6
Senna	<i>Cassia angustifolia</i>	Rhein	Anthraquinone	<i>S. aureus</i>	2.0	
Smooth hydrangea, seven barks	<i>Hydrangea arborescens</i>	?		General	2.3	
Snakeplant	<i>Rivea corymbosa</i>	?		General	1.0	
St. John's wort	<i>Hypericum perforatum</i>	Hypericin, others	Anthraquinone	General	1.7	
Sweet flag, calamus	<i>Acorus calamus</i>	?		Enteric bacteria	0.7	
Tansy	<i>Tanacetum vulgare</i>	Essential oils	Terpenoid	Helminths, bacteria	2.0	
Tarragon	<i>Artemisia dracunculus</i>	Caffeic acids, tannins	Terpenoid	Viruses, helminths	2.5	
			Polyphenols			245

Continued on following page

TABLE 1—Continued

Common name	Scientific name	Compound	Class	Activity ^d	Relative toxicity ^b	Reference(s) ^c
Thyme	<i>Thymus vulgaris</i>	Caffeic acid	Terpenoid	Viruses, bacteria, fungi	2.5	
		Thymol	Phenolic alcohol			
		Tannins	Polyphenols			
Tree bard	<i>Podocarpus nagi</i>	—	Flavones	<i>P. acnes</i> , other gram-positive bacteria		123
		Totanol	Flavonol			
		Nagilactone	Lactone			
Tua-Tua	<i>Jatropha gossypifolia</i>	?		General	0.0	122
Turmeric	<i>Curcuma longa</i>	Curcumin	Terpenoids	Bacteria, protozoa		14
		Turmeric oil				
Valerian	<i>Valeriana officinalis</i>	Essential oil	Terpenoid	General	2.7	
Willow	<i>Salix alba</i>	Salicin	Phenolic glucoside			
		Tannins	Polyphenols			
		Essential oil	Terpenoid			
Wintergreen	<i>Gaultheria procumbens</i>	Tannins	Polyphenols	General	1.0	
Woodruff	<i>Galium odoratum</i>	—	Coumarin	General	3.0	26, 83, 193
Yarrow	<i>Achillea millefolium</i>	?		Viruses, helminths	2.3	24
Yellow dock	<i>Rumex crispus</i>	?		<i>E. coli</i> , <i>Salmonella</i> , <i>Staphylococcus</i>	1.0	

^a Plants with activity only against HIV are listed in Table 6.

^b 0, very safe; 3, very toxic. Data from reference 58.

^c Table is based on data compiled from references 58 and 224; additional references are listed here.

^d "General" denotes activity against multiple types of microorganisms (e.g., bacteria, fungi, and protozoa), and "bacteria" denotes activity against gram-positive and gram-negative bacteria.

isms include enzyme inhibition by the oxidized compounds, possibly through reaction with sulfhydryl groups or through more nonspecific interactions with the proteins (137).

Phenolic compounds possessing a C₃ side chain at a lower level of oxidation and containing no oxygen are classified as essential oils and often cited as antimicrobial as well. Eugenol is a well-characterized representative found in clove oil (Fig. 1). Eugenol is considered bacteriostatic against both fungi (58) and bacteria (224).

Quinones. Quinones are aromatic rings with two ketone substitutions (Fig. 1). They are ubiquitous in nature and are characteristically highly reactive. These compounds, being colored, are responsible for the browning reaction in cut or injured fruits and vegetables and are an intermediate in the melanin synthesis pathway in human skin (194). Their presence in henna gives that material its dyeing properties (69). The switch between diphenol (or hydroquinone) and diketone (or quinone) occurs easily through oxidation and reduction reactions. The individual redox potential of the particular quinone-hydroquinone pair is very important in many biological systems; witness the role of ubiquinone (coenzyme Q) in mammalian electron transport systems. Vitamin K is a complex naphthoquinone. Its antihemorrhagic activity may be related to its ease of oxidation in body tissues (85). Hydroxylated amino acids may be made into quinones in the presence of suitable enzymes, such as a polyphenoloxidase (233). The reaction for the conversion of tyrosine to quinone is shown in Fig. 2.

In addition to providing a source of stable free radicals, quinones are known to complex irreversibly with nucleophilic amino acids in proteins (210), often leading to inactivation of the protein and loss of function. For that reason, the potential range of quinone antimicrobial effects is great. Probable targets in the microbial cell are surface-exposed adhesins, cell

wall polypeptides, and membrane-bound enzymes. Quinones may also render substrates unavailable to the microorganism. As with all plant-derived antimicrobials, the possible toxic effects of quinones must be thoroughly examined.

Kazmi et al. (112) described an anthraquinone from *Cassia italica*, a Pakistani tree, which was bacteriostatic for *Bacillus anthracis*, *Corynebacterium pseudodiphthericum*, and *Pseudomonas aeruginosa* and bactericidal for *Pseudomonas pseudomalliae*. Hypericin, an anthraquinone from St. John's wort (*Hypericum perforatum*), has received much attention in the popular press lately as an antidepressant, and Duke reported in 1985 that it had general antimicrobial properties (58).

Flavones, flavonoids, and flavonols. Flavones are phenolic structures containing one carbonyl group (as opposed to the two carbonyls in quinones) (Fig. 1). The addition of a 3-hydroxyl group yields a flavonol (69). Flavonoids are also hydroxylated phenolic substances but occur as a C₆-C₃ unit linked to an aromatic ring. Since they are known to be synthesized by plants in response to microbial infection (56), it should not be surprising that they have been found in vitro to be effective antimicrobial substances against a wide array of microorganisms. Their activity is probably due to their ability to complex with extracellular and soluble proteins and to complex with bacterial cell walls, as described above for quinones. More lipophilic flavonoids may also disrupt microbial membranes (229).

Catechins, the most reduced form of the C₃ unit in flavonoid compounds, deserve special mention. These flavonoids have been extensively researched due to their occurrence in oolong green teas. It was noticed some time ago that teas exerted antimicrobial activity (227) and that they contain a mixture of catechin compounds. These compounds inhibited in vitro *Vibrio cholerae* O1 (25), *Streptococcus mutans* (23, 185, 186, 228), *Shigella* (235), and other bacteria and microorganisms

TABLE 2. Major classes of antimicrobial compounds from plants

Class	Subclass	Example(s)	Mechanism	Reference(s)	
Phenolics	Simple phenols	Catechol	Substrate deprivation	174	
		Epicatechin	Membrane disruption	226	
	Phenolic acids	Cinnamic acid		66	
		Quinones	Hypericin	Bind to adhesins, complex with cell wall, inactivate enzymes	58, 114
	Flavonoids	Chrysin		Bind to adhesins	175, 182
				Complex with cell wall	
	Flavones	Abyssinone		Inactivate enzymes	32, 219
				Inhibit HIV reverse transcriptase	164
	Flavonols	Totarol		?	122
				Bind to proteins	196, 210
Tannins	Ellagitannin		Bind to adhesins	192	
			Enzyme inhibition	87, 33, 35	
	Coumarins	Warfarin	Interaction with eucaryotic DNA (antiviral activity)	26, 95, 113, 251	
Terpenoids, essential oils		Capsaicin	Membrane disruption	42	
Alkaloids		Berberine	Intercalate into cell wall and/or DNA	15, 34, 73, 94	
		Piperine			
Lectins and polypeptides		Mannose-specific agglutinin	Block viral fusion or adsorption	145, 253	
		Fabatin	Form disulfide bridges		
Polyacetylenes		8S-Heptadeca-2(Z),9(Z)-diene-4,6-diyne-1,8-diol	?	62	

(186, 224). The catechins inactivated cholera toxin in *Vibrio* (25) and inhibited isolated bacterial glucosyltransferases in *S. mutans* (151), possibly due to complexing activities described for quinones above. This latter activity was borne out in in vivo tests of conventional rats. When the rats were fed a diet containing 0.1% tea catechins, fissure caries (caused by *S. mutans*) was reduced by 40% (166).

Flavonoid compounds exhibit inhibitory effects against multiple viruses. Numerous studies have documented the effectiveness of flavonoids such as swertifrancheside (172), glycyrrhizin (from licorice) (242), and chrysin (48) against HIV. More than one study has found that flavone derivatives are inhibitory to respiratory syncytial virus (RSV) (21, 111). Kaul et al. (111) provide a summary of the activities and modes of action of quercetin, naringin, hesperetin, and catechin in in vitro cell culture monolayers. While naringin was not inhibitory to herpes simplex virus type 1 (HSV-1), poliovirus type 1, parainfluenza virus type 3, or RSV, the other three flavonoids were effective in various ways. Hesperetin reduced intracellular replication of all four viruses; catechin inhibited infectivity but not intracellular replication of RSV and HSV-1; and quercetin was universally effective in reducing infectivity. The authors propose that small structural differences in the compounds are critical to their activity and pointed out another advantage of many plant derivatives: their low toxic potential. The average Western daily diet contains approximately 1 g of mixed flavonoids (124); pharmacologically active concentrations are not likely to be harmful to human hosts.

An isoflavone found in a West African legume, alpinumiflavone, prevents schistosomal infection when applied topically (175). Phloretin, found in certain serovars of apples, may have activity against a variety of microorganisms (101). Galangin (3,5,7-trihydroxyflavone), derived from the perennial herb

Helichrysum aureonitens, seems to be a particularly useful compound, since it has shown activity against a wide range of gram-positive bacteria as well as fungi (2) and viruses, in particular HSV-1 and coxsackie B virus type 1 (145).

Delineation of the possible mechanism of action of flavones and flavonoids is hampered by conflicting findings. Flavonoids lacking hydroxyl groups on their β -rings are more active against microorganisms than are those with the $-\text{OH}$ groups (38); this finding supports the idea that their microbial target is the membrane. Lipophilic compounds would be more disruptive of this structure. However, several authors have also found the opposite effect; i.e., the more hydroxylation, the greater the antimicrobial activity (189). This latter finding reflects the similar result for simple phenolics (see above). It is safe to say that there is no clear predictability for the degree of hydroxylation and toxicity to microorganisms.

Tannins. "Tannin" is a general descriptive name for a group of polymeric phenolic substances capable of tanning leather or precipitating gelatin from solution, a property known as astringency. Their molecular weights range from 500 to 3,000 (87), and they are found in almost every plant part: bark, wood, leaves, fruits, and roots (192). They are divided into two groups, hydrolyzable and condensed tannins. Hydrolyzable tannins are based on gallic acid, usually as multiple esters with D-glucose, while the more numerous condensed tannins (often called proanthocyanidins) are derived from flavonoid monomers (Fig. 1). Tannins may be formed by condensations of flavan derivatives which have been transported to woody tissues of plants. Alternatively, tannins may be formed by polymerization of quinone units (76). This group of compounds has received a great deal of attention in recent years, since it was suggested that the consumption of tannin-containing bever-




FIG. 1. Structures of common antimicrobial plant chemicals.

ages, especially green teas and red wines, can cure or prevent a variety of ills (198).

Many human physiological activities, such as stimulation of phagocytic cells, host-mediated tumor activity, and a wide range of anti-infective actions, have been assigned to tannins (87). One of their molecular actions is to complex with proteins through so-called nonspecific forces such as hydrogen bonding

and hydrophobic effects, as well as by covalent bond formation (87, 210). Thus, their mode of antimicrobial action, as described in the section on quinones (see above), may be related to their ability to inactivate microbial adhesins, enzymes, cell envelope transport proteins, etc. They also complex with polysaccharide (247). The antimicrobial significance of this particular activity has not been explored. There is also evidence for

lesser activity against gram-negative bacteria as well as *Candida albicans* (100). Two diterpenes isolated by Batista et al. (23) were found to be more democratic; they worked well against *Staphylococcus aureus*, *V. cholerae*, *P. aeruginosa*, and *Candida* spp. When it was observed that residents of Mali used the bark of a tree called *Ptelopsis suberosa* for the treatment of gastric ulcers, investigators tested terpenoid-containing fractions in 10 rats before and after the rats had ulcers chemically induced. They found that the terpenoids prevented the formation of ulcers and diminished the severity of existent ulcers. Whether this activity was due to antimicrobial action or to protection of the gastric mucosa is not clear (53). Kadota et al. (108) found that trichorabdol A, a diterpene from a Japanese herb, could directly inhibit *H. pylori*.

Alkaloids

Heterocyclic nitrogen compounds are called alkaloids. The first medically useful example of an alkaloid was morphine, isolated in 1805 from the opium poppy *Papaver somniferum* (69); the name morphine comes from the Greek Morpheus, god of dreams. Codeine and heroin are both derivatives of morphine. Diterpenoid alkaloids, commonly isolated from the plants of the Ranunculaceae, or buttercup (107) family (15), are commonly found to have antimicrobial properties (163). Solamargine, a glycoalkaloid from the berries of *Solanum khasianum*, and other alkaloids may be useful against HIV infection (142, 200) as well as intestinal infections associated with AIDS (140). While alkaloids have been found to have microbicidal effects (including against *Giardia* and *Entamoeba* species [78]), the major antidiarrheal effect is probably due to their effects on transit time in the small intestine.

Berberine is an important representative of the alkaloid group. It is potentially effective against trypanosomes (73) and plasmodia (163). The mechanism of action of highly aromatic planar quaternary alkaloids such as berberine and harmaline (93) is attributed to their ability to intercalate with DNA (176).

Lectins and Polypeptides

Peptides which are inhibitory to microorganisms were first reported in 1942 (19). They are often positively charged and contain disulfide bonds (253). Their mechanism of action may be the formation of ion channels in the microbial membrane (222, 253) or competitive inhibition of adhesion of microbial proteins to host polysaccharide receptors (202). Recent interest has been focused mostly on studying anti-HIV peptides and lectins, but the inhibition of bacteria and fungi by these macromolecules, such as that from the herbaceous *Amaranthus*, has long been known (50).

Thionins are peptides commonly found in barley and wheat and consist of 47 amino acid residues (45, 143). They are toxic to yeasts and gram-negative and gram-positive bacteria (65). Thionins AX1 and AX2 from sugar beet are active against fungi but not bacteria (118). Fabatin, a newly identified 47-residue peptide from fava beans, appears to be structurally related to γ -thionins from grains and inhibits *E. coli*, *P. aeruginosa*, and *Enterococcus hirae* but not *Candida* or *Saccharomyces* (253).

The larger lectin molecules, which include mannose-specific lectins from several plants (20), MAP30 from bitter melon (128), GAP31 from *Gelonium multiflorum* (28), and jacalin (64), are inhibitory to viral proliferation (HIV, cytomegalovirus), probably by inhibiting viral interaction with critical host cell components. It is worth emphasizing that molecules and compounds such as these whose mode of action may be to inhibit adhesion will not be detected by using most general

plant antimicrobial screening protocols, even with the bioassay-guided fractionation procedures (131, 181) used by natural-products chemists (see below). It is an area of ethnopharmacology which deserves attention, so that initial screens of potentially pharmacologically active plants (described in references 25, 43, and 238) may be made more useful.

Mixtures

The chewing stick is widely used in African countries as an oral hygiene aid (in place of a toothbrush) (156). Chewing sticks come from different species of plants, and within one stick the chemically active component may be heterogeneous (5). Crude extracts of one species used for this purpose, *Serindeia werneckii*, inhibited the periodontal pathogens *Porphyromonas gingivalis* and *Bacteroides melaninogenicus* in vitro (183). The active component of the Nigerian chewing stick (*Fagara zanthoxyloides*) was found to consist of various alkaloids (157). Whether these compounds, long utilized in developing countries, might find use in the Western world is not yet known.

Papaya (*Carica papaya*) yields a milky sap, often called a latex, which is a complex mixture of chemicals. Chief among them is papain, a well-known proteolytic enzyme (162). An alkaloid, carpaine, is also present (34). Terpenoids are also present and may contribute to its antimicrobial properties (224). Osato et al. (168) found the latex to be bacteriostatic to *B. subtilis*, *Enterobacter cloacae*, *E. coli*, *Salmonella typhi*, *Staphylococcus aureus*, and *Proteus vulgaris*.

Ayurveda is a type of healing craft practiced in India but not unknown in the United States. Ayurvedic practitioners rely on plant extracts, both "pure" single-plant preparations and mixed formulations. The preparations have lyrical names, such as Ashwagandha (*Withania somnifera* root) (54), Cauvery 100 (a mixture) (133), and Livo-vet (125). These preparations are used to treat animals as well as humans (125). In addition to their antimicrobial activities, they have been found to have antidiarrheal (134), immunomodulatory (54, 133), anticancer (59), and psychotropic (201) properties. In vivo studies of Abana, an Ayurvedic formulation, found a slight reduction in experimentally induced cardiac arrhythmias in dogs (75). Two microorganisms against which Ayurvedic preparations have activity are *Aspergillus* spp. (54) and *Propionibacterium acnes* (170). (The aspergillosis study was performed with mice in vivo, and it is therefore impossible to determine whether the effects are due to the stimulation of macrophage activity in the whole animal rather than to direct antimicrobial effects.)

The toxicity of Ayurvedic preparations has been the subject of some speculation, especially since some of them include metals. Prpic-Majic et al. identified high levels of lead in the blood of adult volunteers who had self-medicated with Ayurvedic medicines (178).

Propolis is a crude extract of the balsam of various trees; it is often called bee glue, since honeybees gather it from the trees. Its chemical composition is very complex: like the latexes described above, terpenoids are present, as well as flavonoids, benzoic acids and esters, and substituted phenolic acids and esters (9). Synthetic cinnamic acids, identical to those from propolis, were found to inhibit hemagglutination activity of influenza virus (199). Amoros et al. found that propolis was active against an acyclovir-resistant mutant of HSV-1, adenovirus type 2, vesicular stomatitis virus, and poliovirus (9). Mixtures of chemicals, such as are found in latex and propolis, may act synergistically. While the flavone and flavonol components were active in isolation against HSV-1, multiple flavonoids incubated simultaneously with the virus were more effective

TABLE 3. Solvents used for active component extraction^a

Water	Ethanol	Methanol	Chloroform	Dichloromethanol	Ether	Acetone
Anthocyanins (111) Starches Tannins (192) Saponins (53) Terpenoids Polypeptides Lectins	Tannins (204) Polyphenols (151) Polyacetylenes (29, 62) Flavonol (29, 100) Terpenoids (82) Sterols (53) Alkaloids (103) Propolis	Anthocyanins Terpenoids (221) Saponins Tannins (221) Xanthoxyllines Totarol (121) Quassinoids (115) Lactones (180) Flavones (189, 219) Phenones (174) Polyphenols (235)	Terpenoids (18) Flavonoids (175)	Terpenoids (144)	Alkaloids Terpenoids Coumarins Fatty acids	Flavonols (2)

^a Compounds in bold are commonly obtained only in one solvent. References are given in parentheses.

than single chemicals, a possible explanation of why propolis is more effective than its individual compounds (10). Of course, mixtures are more likely to contain toxic constituents, and they must be thoroughly investigated and standardized before approved for use on a large-scale basis in the West.

Other Compounds

Many phytochemicals not mentioned above have been found to exert antimicrobial properties. This review has attempted to focus on reports of chemicals which are found in multiple instances to be active. It should be mentioned, however, that there are reports of antimicrobial properties associated with polyamines (in particular spermidine) (70), isothiocyanates (57, 104), thiosulfates (215), and glucosides (149, 184).

Polyacetylenes deserve special mention. Estevez-Braun et al. isolated a C₁₇ polyacetylene compound from *Bupleurum salicifolium*, a plant native to the Canary Islands. The compound, 8*S*-heptadeca-2(*Z*),9(*Z*)-diene-4,6-diyne-1,8-diol, was inhibitory to *S. aureus* and *B. subtilis* but not to gram-negative bacteria or yeasts (62). Acetylene compounds and flavonoids from plants traditionally used in Brazil for treatment of malaria fever and liver disorders have also been associated with anti-malarial activity (29).

Much has been written about the antimicrobial effects of cranberry juice. Historically, women have been told to drink the juice in order to prevent and even cure urinary tract infections. In the early 1990s, researchers found that the monosaccharide fructose present in cranberry and blueberry juices competitively inhibited the adsorption of pathogenic *E. coli* to urinary tract epithelial cells, acting as an analogue for mannose (252). Clinical studies have borne out the protective effects of cranberry juice (17). Many fruits contain fructose, however, and researchers are now seeking a second active compound from cranberry juice which contributes to the antimicrobial properties of this juice (252).

EXPERIMENTAL APPROACHES

Extraction Methods

Advice abounds for the amateur herbalist on how to prepare healing compounds from plants and herbs. Water is almost universally the solvent used to extract activity. At home, dried plants can be ingested as teas (plants steeped in hot water) or, rarely, tinctures (plants in alcoholic solutions) or inhaled via steam from boiling suspensions of the parts. Dried plant parts can be added to oils or petroleum jelly and applied externally. Poultices can also be made from concentrated teas or tinctures (30, 224).

Scientific analysis of plant components follows a logical pathway. Plants are collected either randomly or by following leads supplied by local healers in geographical areas where the plants are found (135). Initial screenings of plants for possible antimicrobial activities typically begin by using crude aqueous or alcohol extractions and can be followed by various organic extraction methods. Since nearly all of the identified components from plants active against microorganisms are aromatic or saturated organic compounds, they are most often obtained through initial ethanol or methanol extraction. In fact, many studies avoid the use of aqueous fractionation altogether. The exceptional water-soluble compounds, such as polysaccharides (e.g., starch) and polypeptides, including fabatin (253) and various lectins, are commonly more effective as inhibitors of pathogen (usually virus) adsorption and would not be identified in the screening techniques commonly used. Occasionally tannins and terpenoids will be found in the aqueous phase, but they are more often obtained by treatment with less polar solvents. Table 3 lists examples of extraction solvents and the resultant active fractions reported in recent studies. Compounds which, according to the literature, partition exclusively in particular solvents are indicated in boldface type in the table.

Any part of the plant may contain active components. For instance, the roots of ginseng plants contain the active saponins and essential oils, while eucalyptus leaves are harvested for their essential oils and tannins. Some trees, such as the balsam poplar, yield useful substances in their bark, leaves, and shoots (224).

For alcoholic extractions, plant parts are dried, ground to a fine texture, and then soaked in methanol or ethanol for extended periods. The slurry is then filtered and washed, after which it may be dried under reduced pressure and redissolved in the alcohol to a determined concentration. When water is used for extractions, plants are generally soaked in distilled water, blotted dry, made into a slurry through blending, and then strained or filtered. The filtrate can be centrifuged (approximately 20,000 × *g*, for 30 min) multiple times for clarification (42, 221). Crude products can then be used in disc diffusion and broth dilution assays to test for antifungal and antibacterial properties and in a variety of assays to screen for antiviral activity, as described below.

Natural-products chemists further purify active chemicals from crude extracts by a variety of methods. Petalostemumol, a flavanol from purple prairie clover, was obtained from the ethanol extract by partitioning between ethyl acetate and water, followed by partitioning between *n*-hexane and 10% methanol. The methanol fraction was chromatographed and eluted with toluene (100). Terpenoid lactones have been obtained by

TABLE 4. Comparison of extractants on different parameters based on a five-point scale (1 to 5) and with different weights allocated to the different parameters^a

Parameter	Weight ^b	Acetone		Ethanol		Methanol		MCW ^c		Methylene dichloride		Water	
		A ^d	C ^e	A	C	A	C	A	C	A	C	A	C
Quantity extracted	3	6	3	9	6	12	12	12	12	3	3	9	9
Rate of extraction	3	12	15	12	12	12	12	12	12	15	15	9	9
No. of compounds extracted	5	20	20	10	15	15	20	10	15	10	15	5	5
No. of inhibitors extracted	5	20	20	0	10	20	15	20	20	20	15	0	0
Toxicity in bioassay	4	16	16	8	8	0	0	8	8	8	8	16	16
Ease of removal	5	20	20	5	5	10	10	10	10	20	20	0	0
Hazardous to use	2	8	8	8	8	2	2	6	6	6	6	8	8
Total		102	102	52	64	71	71	78	83	79	79	47	47

^a Reprinted from reference 61 with permission of the publisher.

^b Five-point scale (5, best; 1, worst), based on Eloff's (61) judgment of the importance of the parameter.

^c MCW, methanol-chloroform-water.

^d Results for *Anthocleista grandiflora*.

^e Results for *Combretum erythrophyllum*.

successive extractions of dried bark with hexane, CHCl₃, and methanol, with activity concentrating in the CHCl₃ fraction (180). The chemical structures of the purified material can then be analyzed. Techniques for further chemical analysis include chromatography, bioautography, radioimmunoassay, various methods of structure identification, and newer tools such as fast atom bombardment mass spectrometry, tandem mass spectrometry (181), high-performance liquid chromatography, capillary zone electrophoresis, nuclear magnetic resonance spectroscopy, and X-ray crystallography (25).

Recently, Eloff (61) examined a variety of extractants for their ability to solubilize antimicrobials from plants, as well as other factors such as their relative ranking as biohazards and the ease of removal of solvent from the fraction. The focus of the study was to provide a more standardized extraction method for the wide variety of researchers working in diverse settings. Although it is not one of the more frequently used extractants in studies published to date, acetone received the highest overall rating. In fact, in a review of 48 articles describing the screening of plant extracts for antimicrobial properties in the most recent years of the *Journal of Natural Products*, the *Journal of Ethnopharmacology*, and the *International Journal of Pharmacognosy*, only one study used acetone as an extractant. Of the solvents listed in Table 3, which are reported in the recent literature with the highest frequency, Eloff ranked them in the order methylene dichloride, methanol, ethanol, and water. Table 4 displays the breakdown of scores for the various solvents studied. The row indicating the number of inhibitors extracted with each solvent points to two implications: first, that most active components are not water soluble, supporting the data reported in Table 3, and second, that the most commonly used solvents (ethanol and methanol, both used as initial extractants in approximately 35% of the studies appearing in the recent literature) may not demonstrate the greatest sensitivity in yielding antimicrobial chemicals on an initial screening. This disparity should be examined as the search for new antimicrobials intensifies.

Efficacy

In vitro experiments. (i) Bacteria and fungi. Initial screening of potential antibacterial and antifungal compounds from plants may be performed with pure substances (2, 23, 117) or crude extracts (73, 203, 182). The methods used for the two types of organisms are similar. The two most commonly used screens to determine antimicrobial susceptibility are the broth

dilution assay (18, 89, 219) and the disc or agar well diffusion assay (153); clinical microbiologists are very familiar with these assays. Adaptations such as the agar overlay method (138) may also be used. In some cases, the inoculated plates or tubes are exposed to UV light (221) to screen for the presence of light-sensitizing photochemicals. Other variations of these methods are also used. For instance, to test the effects of extracts on invasive *Shigella* species, noncytotoxic concentrations of the extracts can be added to Vero cell cultures exposed to a *Shigella* inoculum (235). The decrease in cytopathic effect in the presence of the plant extract is then measured.

In addition to these assays, antifungal phytochemicals can be analyzed by a spore germination assay. Samples of plant extracts or pure compounds can be added to fungal spores collected from solid cultures, placed on glass slides, and incubated at an appropriate temperature (usually 25°C) for 24 h. Slides are then fixed in lactophenol-cotton blue and observed microscopically for spore germination (179).

Of course, after initial screening of phytochemicals, more detailed studies of their antibiotic effects should be conducted. At this stage, more specific media can be used and MICs can be effectively compared to those of a wider range of currently used antibiotics. The investigation of plant extracts effective against methicillin-resistant *S. aureus* (190, 229) provides an example of prospecting for new compounds which may be particularly effective against infections that are currently difficult to treat. Sato et al. (190) examined the activity of three extracts from the fruiting bodies of the tree *Terminalia chebula* RETS against methicillin-sensitive and methicillin-resistant *S. aureus* as well as 12 other gram-negative and gram-positive bacteria. They found that gallic acid derivatives were more effective against both types of *S. aureus* than they were against other species.

(ii) Viruses. Several methods are available to detect either virucidal or inhibitory (antiviral) plant activity. Investigators can look for cytopathic effects or plaque formation (1, 111) or for transformation or proliferative effects on cell lines (48). Viral replication may be assayed by detection of viral products such as DNA, RNA, or polypeptides (127, 171). Vlietinck and Vanden Berghe (238) have noted that the method for assaying antiviral substances used in various laboratories is not standardized, and therefore the results are often not comparable to one another. These authors also point out that researchers must distinguish between merely toxic effects of agents on host

cells and true antiviral properties of the plant extracts. Table 5 lists various in vitro antiviral screening assays.

It should be mentioned here that antiviral assays often screen for active substances which prevent adsorption of the microorganism to host cells; this activity is overlooked in screening procedures for antibacterial and antifungal substances. However, the large body of literature accumulating on antiadhesive approaches to bacterial infections (109) suggests that phytochemicals should be assessed for this class of action in addition to their killing and inhibitory activities.

(iii) Protozoa and helminths. Screening plant extracts for activity against protozoa and helminths can be more complicated than screening for activity against bacteria, fungi, or viruses. Growing the organisms is often more difficult, and fewer organisms are obtained. Assays are generally specific for the microorganism. For instance, Freiburghaus et al. (73) used two methods to assay compounds for effectiveness against *Trypanosoma brucei*. First, 66-h tissue cultures of the trypanosomes were exposed to various concentrations of extracts, and then the MICs were determined with an inverted microscope. (In this case, the MIC was defined as the lowest concentration of extract which completely inhibited the growth of the trypanosomes.) Second, these authors used a fluorescence assay of trypanosome viability in microtiter wells.

There is great interest in plant chemicals which may have anti-infective properties for *Plasmodium* species. One method for testing extracts which may be toxic to plasmodia uses radiolabelled microorganisms in an erythrocyte infection assay with microtiter dishes (72). Parasitized erythrocytes are incubated with and without test substances, and the numbers of *Plasmodium* organisms after the incubation period are quantitated. Another laboratory directly measures decreases in the growth of *Plasmodium* under the influence of the extracts (29).

In vivo testing of phytochemicals. Two prime examples of animal studies from the recent literature are the descriptions of the effects of oolong tea polyphenols on dental caries in rats (186) and cholera in mice (226) (discussed above). Another bacterial infection model, *P. aeruginosa* lung infection of athymic rats, was used to test subcutaneous administrations of ginseng. Treated rats had decreased lung pathology and bacterial load, although enhanced humoral immunity was thought to be at least partially responsible (207).

Perhaps because of the long-standing interest in alternative pharmaceuticals for use in developing and tropical regions of the world, many of the in vivo studies to date have examined the antiprotozoal effectiveness of plant extracts (55). Two recent studies with rats used experimental *Entamoeba* infections to test the effectiveness of a crude Indian herbal extract (205) and of an ethanolic extract of *Piper longum* fruits, probably containing the alkaloid piperine (78). After applying the chloroform extract of the legume *Milletia thonningii* topically to the skin of mice, Perrett et al. (175) found that it prevented the establishment of subsequent *Schistosoma mansoni* infection. There are also animal studies of the activity of various extracts of the bark of *Pteleopsis suberosa* against ulcers (53), of papaya latex against infections with the helminth *Heligmosomoides polygyrus* (191), and of *Santolina chamaecyparissus* essential oil against candidal infections (213).

An Ayurvedic formulation, Cauvery 100, was administered orally to rats either before or after diarrhea was induced by castor oil. While this trial did not test antimicrobial activities of the preparation but, rather, its effect on gut motility and ion loss in the sick animals, it showed that Cauvery 100 had both a protective and a curative effect in rats (134). A similar study with similar results was performed on rats with chemically induced ulcers (133). The Ayurvedic Ashwagandha was found

TABLE 5. In vitro antiviral screening assays^a

Determination of viral infectivity under two conditions:	
I.	In cultured cells during virus multiplication in the presence of a single compound (A-S) or a mixture of compounds, e.g., plant extracts (A-M).
II.	After extracellular incubation with a single compound (V-S) or a mixture of compounds (V-M).
Plaque inhibition assay (only for viruses which form plaques in suitable cell systems)	
	Titer determination of a limited number of viruses in the presence of a nontoxic dose of the test substance
	Applicability: A-S
Plaque reduction assay (only for viruses which form plaques in suitable cell systems)	
	Titer determination of residual virus infectivity after extracellular action of test substance(s) (see reference for comments on cytotoxicity)
	Applicability: V-S, V-M
Inhibition of virus-induced cytopathic effect (for viruses that induce cytopathic effect but do not readily form plaques in cell cultures)	
	Determination of virus-induced cytopathic effect in cell monolayers cultured in liquid medium, infected with a limited dose of virus, and treated with a nontoxic dose of the test substance(s)
	Applicability: A-S, A-M
Virus yield reduction assay	
	Determination of the virus yield in tissues cultures infected with a given amount of virus and treated with a nontoxic dose of the test substance(s)
	Virus titer determination carried out after virus multiplication by the plaque test or the 50% tissue culture infective dose end-point test
	Applicability: A-S, A-M
End-point titer determination technique	
	Determination of virus titer reduction in the presence of twofold dilutions of test compound(s)
	This method especially designed for the antiviral screening of crude extracts (238)
	Applicability: A-S, A-M
Assays based on measurement of specialized functions and viral products (for viruses that do not induce cytopathic effects or form plaques in cell cultures)	
	Determination of virus specific parameters, e.g., hemagglutination and hemadsorption tests (myxoviruses), inhibition of cell transformation (Epstein-Barr virus), immunological tests detecting antiviral antigens in cell cultures (Epstein-Barr virus, HIV, HSV, cytomegalovirus)
	Applicability: A-S, A-M, V-S, V-M
	Reduction or inhibition of the synthesis of virus specific polypeptides in infected cell cultures, e.g., viral nucleic acids, determination of the uptake of radioactive isotope-labelled precursors or viral genome copy numbers
	Applicability: A-S, A-M, V-S, V-M

^a Reprinted from reference 238 with permission of the publisher.

to prolong the survival period of BALB/c mice infected with *Aspergillus*, an effect believed to be mediated by increased macrophage function (54).

Clinical trials in humans. Of course, plants have been used for centuries to treat infections and other illnesses in humans in aboriginal groups, but controlled clinical studies are scarce. In some cases, traditional healers working together with trained scientists have begun keeping records of the safety and effectiveness of phytochemical treatments, but these are generally uncontrolled and unrandomized studies. Examples of

TABLE 6. Compounds with anti-HIV activity

Target	Compound	Class	Plant	Reference(s)
Reverse transcriptase	Ellagitannin	Tannin	— ^a	155
	Hydroxymaprounic acid, hydroxybenzoate	Terpenoids	<i>Maprounea africana</i>	173
	Betulinic acid, platanic acid	Terpenoids	<i>Syzigium claviflorum</i>	74
	Catechin	Polyphenol	—	147
	Faicalcin, quercetin, myricetin, baicalin	Flavonoids	<i>Quercus rubra</i> , others	165, 208, 216
	Nigranoic acid	Terpenoids	<i>Schisandra sphaerandra</i>	212
	Amentoflavone, scutellarein, others	Flavonoids, flavones	—	96, 146, 164, 165, 208, 217, 218
	Benzophenanthridine	Alkaloid	—	200, 217
	Protoberberine	Alkaloid	—	217
	Psychotrines	Alkaloids	Ipecac (<i>Cephaelis ipecacuanha</i>)	217
	Michellamine B	Alkaloid	<i>Ancistrocladus korupensis</i>	142
	Suksdorfii	Coumarins	<i>Lomatium suksdorfii</i>	126
	Coriandrin	Coumarin	Coriander (<i>Coriandrum sativum</i>)	99
	Caffeic acid	Tannin	<i>Hyssop officinalis</i>	119
	Cornusin, others	Condensed and hydrolyzable tannins	<i>Cornus officinalis</i> and others	111
	Swertifrancheside	Flavonoid	<i>Swertia franchetiana</i>	172
	Salaspermic acid	Flavonoid	<i>Trypterygium wilfordii</i>	40
	Glycyrrhizin	Flavonoid	Licorice (<i>Glycyrrhiza rhiza</i>)	102, 242
	—	Protein	Cactus (<i>Opuntia streptacantha</i>)	3
	Methyl nordihydroguaiaretic acid	Lignan	Many trees	80
	Thuja polysaccharide	Polysaccharide	<i>Thuja occidentalis</i>	160
		Lignin-polysaccharide complexes	Japanese white pine (<i>Pinus parviflora</i>)	126
	Integrase	—	Flavonoids, flavones	—
MAP30, GAP31, DAP 32, DAP 30		Proteins	<i>Momordica charantia</i> , <i>Gelonium multiflorum</i>	97, 128–130
—		Lectins	—	20
Caffeic acid		Terpenoid	—	68
Curcumin		Terpenoid	Turmeric (<i>Curcuma longa</i>)	139
Quercetin		Flavonoid	<i>Quercus rubra</i>	67
Protease	Carnosolic acid	Terpenoid	Rosemary (<i>Rosmarinus officinalis</i>)	171
	Ursolic acid	Terpenoids	<i>Geum japonicum</i>	246
		Flavonoids, flavones		32, 48, 67
Adsorption	Mannose-specific lectins	Lectins	Snowdrop (<i>Galanthus</i>), daffodil (<i>Narcissus</i>), amaryllis (<i>Amaryllis</i>), <i>Gerardia</i>	148
	Schumannificine 1	Alkaloid	<i>Schumanniphyton magnificum</i>	94
	Prunellin	Polysaccharide	<i>Prunella</i>	249
Viral fusion	Mannose- and <i>N</i> -acetylglucosamine-specific lectins	Lectins	<i>Cymbidium</i> hybrid, <i>Epiactis helleborine</i> , <i>Listeria ovata</i> , <i>Urtica dioica</i>	20
Syncytium formation	Propolis	Mixture	Various trees	9, 52
	Michellamine B	Alkaloid	<i>Ancistrocladus korupensis</i>	142
	MAR-10	Polysaccharide	<i>Hyssop officinalis</i>	81
Interference with cellular factors	Chrysin	Flavone	<i>Chrysanthemum morifolium</i>	48
	29,000-mol-wt protein	Protein	Pokeweed (<i>Phytolacca</i>)	41
	Hypericin	Anthraquinone	St. John's wort (<i>Hypericum</i>)	41, 99
	Camptothecin	Alkaloid	?	169
	Trichosanthin, momorcharins	Ribosome-inactivating proteins	Cucurbitaceae family	141
Unknown	—	Sulfated polysaccharide	<i>Prunella vulgaris</i>	214
	—	Sulfonated polysaccharide	<i>Viola yedoensis</i>	154
	Jacalin	Lectin	?	46, 64
	Zingibroside R-1	Terpenoid	<i>Panax zingiberensis</i> Wu et Feng	86
	Thiarubrines	Terpenoids	Asteraceae	98
	Chrysin	Flavonoid	<i>Chrysanthemum morifolium</i>	96
	—	Essential oil/terpenoids	<i>Houttuynia cordata</i>	88

^a —, unspecified.

peer-reviewed trials of the use of phytochemicals in humans are briefly reported here.

A cross-sectional epidemiological study (thus, not a randomized trial) of the effectiveness of chewing sticks versus toothbrushes in promoting oral hygiene was conducted in West Africa. The study's authors found a reduced effectiveness in chewing-stick users compared to toothbrush users and concluded that the antimicrobial chemicals known to be present in the sticks added no oral health benefit (156). Also, regarding oral health, mouthrinses containing various antimicrobials have been evaluated in humans (reviewed in reference 240). Mouthrinses containing phytochemicals were not found to be as effective in decreasing plaque or clinical gingivitis as were Listerine or chlorhexidine.

To date, few randomized clinical trials of plant antimicrobials have been performed. Giron et al. (79) compared an extract of *Solanum nigrescens* with nystatin, both given as intravaginal suppositories, in women with confirmed *C. albicans* vaginitis. The extract proved to be as efficacious as nystatin. In 1996 the long-recognized effects of cranberry juice on urinary tract infections were studied in a group of elderly women. No data are provided by the author about clinical outcomes, but treated women had fewer bacteria in their urine than did the untreated group (17).

Four different Ayurvedic formulations were tested against a placebo for their effectiveness against acne vulgaris. One of the formulations, Sunder Vati, produced a significantly greater reduction in lesion count than did the other three and the placebo. No toxic side effects were observed (170). Two proprietary compounds derived from tropical plants, Provir, for the treatment of respiratory viral infections, and Virend, a topical antiherpes agent, were tested in clinical trials in 1994 (114). Since then, the efficacy and safety of Virend have been established in phase II studies (167).

PLANTS WITH EFFICACY AGAINST HUMAN IMMUNODEFICIENCY VIRUS

Effective therapies for HIV infection are being sought far and wide, in the natural world as well as in laboratories. As one example of in vivo anti-HIV studies, infection in mice has been studied. Glycyrrhizin, found in *Glycyrrhiza* plants (the source of licorice), extended the life of the retrovirus-infected mice from 14 to 17 weeks (242). A crude extract of the cactus *Opuntia streptacantha* had marked antiviral effects in vitro, and toxicity studies performed in mice, horses, and humans found the extract to be safe (3).

The scope of studies of anti-HIV plant extracts is too broad to handle in detail in the present review, but Table 6 summarizes many of the compounds studied to date as well as their purported targets of action. The interested reader is referred to several useful reviews (36, 155, 172, 217).

COMMERCIAL AVAILABILITY AND SAFETY OF COMPOUNDS

A wide variety of plant extracts, mixtures, and single plant compounds are available in the United States without a prescription through health food stores and vitamin retailers. For example, preparations of flavones (brand names Flavons 500 and Citrus Bioflavonoids) are sold by supplement suppliers. Capsaicin appears in a topical antiarthritis cream available in mainstream grocery and drug stores. The over-the-counter preparations are relatively unregulated. This results in preparations of uncertain and variable purity.

Many stores also sell whole dried plants, which have been

found occasionally to be misidentified, with potentially disastrous consequences. Obviously, many plants can also be grown by the consumer in home gardens. In 1994, Congress passed the Dietary Supplement Health Education Act, which required the Food and Drug Administration to develop labeling requirements specifically designed for products containing ingredients such as vitamins, minerals, and herbs intended to supplement the diet. The new rules, issued in late 1997, require products to be labeled as a dietary supplement and carry a "Supplement Facts" panel with information similar to the "Nutrition Facts" panels appearing on processed foods. The rules also require that products containing botanical ingredients specify the part of the plant used (71).

The Food and Drug Administration regulates dietary supplements as foods, as long as no drug claims are made for them (63). Other organizations overseeing herbals and botanicals are the American Dietetic Association's Commission on Dietary Supplement Labels (11) and the Food and Nutrition Science Alliance (12).

The first investigational new drug application for herbal pharmaceuticals, available by prescription, was submitted in 1997 (13). Whether this is the beginning of a substantial "rush to market" is unclear. Some analysts suggest that pharmaceutical companies will not invest in drug testing for botanicals since it is still unclear whether proprietary claims can be made (49). Richard van den Broek, industry analyst, concurs, saying that most novel antibiotic research is being conducted instead by biotechnology companies (7).

Plants and/or plant components which are currently available to consumers as supplements through commercial outlets are among those listed in Table 1. Rough toxicity rankings are also provided. It has long been recognized that possibly therapeutic foodstuffs may also contain substances which act as poisons in humans (132). Clinical scientists and practitioners are best advised to be aware of the widespread self-medication with these products and to consider their effects on patients.

CONCLUSIONS AND FUTURE DIRECTIONS

Scientists from divergent fields are investigating plants anew with an eye to their antimicrobial usefulness. A sense of urgency accompanies the search as the pace of species extinction continues. Laboratories of the world have found literally thousands of phytochemicals which have inhibitory effects on all types of microorganisms in vitro. More of these compounds should be subjected to animal and human studies to determine their effectiveness in whole-organism systems, including in particular toxicity studies as well as an examination of their effects on beneficial normal microbiota. It would be advantageous to standardize methods of extraction and in vitro testing so that the search could be more systematic and interpretation of results would be facilitated. Also, alternative mechanisms of infection prevention and treatment should be included in initial activity screenings. Disruption of adhesion is one example of an anti-infection activity not commonly screened for currently. Attention to these issues could usher in a badly needed new era of chemotherapeutic treatment of infection by using plant-derived principles.

ACKNOWLEDGMENT

I gratefully acknowledge the technical assistance of Emily A. Horst.

REFERENCES

1. Abou-Karam, M., and W. T. Shier. 1990. A simplified plaque reduction assay for antiviral agents from plants. Demonstration of frequent occurrence of antiviral activity in higher plants. *J. Nat. Prod.* 53:340-344.

2. Afolayan, A. J., and J. J. M. Meyer. 1997. The antimicrobial activity of 3,5,7-trihydroxyflavone isolated from the shoots of *Helichrysum aureonitens*. *J. Ethnopharmacol.* **57**:177–181.
3. Ahmad, A., J. Davies, S. Randall, and G. R. B. Skinner. 1996. Antiviral properties of extract of *Opuntia streptacantha*. *Antiviral Res.* **30**:75–85.
4. Ahmed, A. A., A. A. Mahmoud, H. J. Williams, A. I. Scott, J. H. Reibenspies, and T. J. Mabry. 1993. New sesquiterpene α -methylene lactones from the Egyptian plant *Jasonia candicans*. *J. Nat. Prod.* **56**:1276–1280.
5. Akpata, E. S., and E. O. Akinrimisi. 1977. Antibacterial activity of extracts from some African chewing sticks. *Oral Surg. Oral Med. Oral Pathol.* **44**:717–722.
6. Ali-Shtayeh, M. S., M. A. Al-Nuri, R. M. R. Yaghmour, and Y. R. Faidi. 1997. Antimicrobial activity of *Micromeria nervosa* from the Palestinian area. *J. Ethnopharmacol.* **58**:143–147.
7. Alper, J. 1998. Effort to combat microbial resistance lags. *ASM News* **64**:440–441.
8. Amaral, J. A., A. Ekins, S. R. Richards, and R. Knowles. 1998. Effect of selected monoterpenes on methane oxidation, denitrification, and aerobic metabolism by bacteria in pure culture. *Appl. Environ. Microbiol.* **64**:520–525.
9. Amoros, M., F. Sauvager, L. Girre, and M. Cormier. 1992. In vitro antiviral activity of propolis. *Apidologie* **23**:231–240.
10. Amoros, M., C. M. O. Simoes, and L. Girre. 1992. Synergistic effect of flavones and flavonols against herpes simplex virus type 1 in cell culture. Comparison with the antiviral activity of propolis. *J. Nat. Prod.* **55**:1732–1740.
11. Anonymous. 1998. Commission on Dietary Supplement Labels issues final report. *J. Am. Diet. Assoc.* **98**:270.
12. Anonymous. 1997. FANSA releases statement about dietary supplement labeling. *J. Am. Diet. Assoc.* **97**:728–729.
13. Anonymous. 1997. First IND submitted with FDA for an herbal pharmaceutical. *AIDS Weekly* Plus 18.
14. Apisariyakul, A., N. Vanittanakom, and D. Buddhasukh. 1995. Antifungal activity of turmeric oil extracted from *Curcuma longa* (Zingiberaceae). *J. Ethnopharmacol.* **49**:163–169.
15. Atta-ur-Rahman and M. I. Choudhary. 1995. Diterpenoid and steroidal alkaloids. *Nat. Prod. Rep.* **12**:361–379.
16. Aureli, P., A. Costantini, and S. Zolea. 1992. Antimicrobial activity of some plant essential oils against *Listeria monocytogenes*. *J. Food Prot.* **55**:344–348.
17. Avorn, J. 1996. The effect of cranberry juice on the presence of bacteria and white blood cells in the urine of elderly women. What is the role of bacterial adhesion? *Adv. Exp. Med. Biol.* **408**:185–186.
18. Ayafor, J. F., M. H. K. Tchuendem, and B. Nyasse. 1994. Novel bioactive diterpenoids from *Aframomum aulacocarpos*. *J. Nat. Prod.* **57**:917–923.
19. Balls, A. K., W. S. Hale, and T. H. Harris. 1942. A crystalline protein obtained from a lipoprotein of wheat flour. *Cereal Chem.* **19**:279–288.
20. Balzarini, J., D. Schols, J. Neyts, E. Van Damme, W. Peumans, and E. De Clercq. 1991. α -(1,3)- and α -(1,6)-D-mannose-specific plant lectins are markedly inhibitory to human immunodeficiency virus and cytomegalovirus infections in vitro. *Antimicrob. Agents Chemother.* **35**:410–416.
21. Barnard, D. L., J. H. Huffman, L. R. Meyerson, and R. W. Sidwell. 1993. Mode of inhibition of respiratory syncytial virus by a plant flavonoid. *Chemotherapy* **39**:212–217.
22. Barre, J. T., B. F. Bowden, J. C. Coll, J. Jesus, V. E. Fuente, G. C. Janairo, and C. Y. Ragasa. 1997. A bioactive triterpene from *Lantana camara*. *Phytochemistry* **45**:321–324.
23. Batista, O., A. Duarte, J. Nascimento, and M. F. Simones. 1994. Structure and antimicrobial activity of diterpenes from the roots of *Plectranthus hereroensis*. *J. Nat. Prod.* **57**:858–861.
24. Berkada, B. 1978. Preliminary report on warfarin for the treatment of herpes simplex. *J. Irish Coll. Phys. Surg.* **22**(Suppl.):56.
25. Borris, R. P. 1996. Natural products research: perspectives from a major pharmaceutical company. *J. Ethnopharmacol.* **51**:29–38.
26. Bose, P. K. 1958. On some biochemical properties of natural coumarins. *J. Indian Chem. Soc.* **58**:367–375.
27. Bosland, P. W. 1994. Chiles: history, cultivation and uses, p. 347–366. *In* G. Charalambous (ed.), *Spices, herbs and edible fungi*. Elsevier, New York, N.Y.
28. Bourinbaier, A. S., and S. Lee Huang. 1996. The activity of plant-derived antiretroviral proteins MAP30 and GAP31 against herpes simplex virus infection in vitro. *Biochem. Biophys. Res. Commun.* **219**:923–929.
29. Brandao, M. G. L., A. U. Krettli, L. S. R. Soares, C. G. C. Nery, and H. C. Marinuzzi. 1997. Antimalarial activity of extracts and fractions from *Bidens pilosa* and other *Bidens* species (Asteraceae) correlated with the presence of acetyle and flavonoid compounds. *J. Ethnopharmacol.* **57**:131–138.
30. Brantner, A., and E. Grein. 1994. Antibacterial activity of plant extracts used externally in traditional medicine. *J. Ethnopharmacol.* **44**:35–40.
31. Brantner, A., Z. Males, S. Pepeljnjak, and A. Antolic. 1996. Antimicrobial activity of *Paliurus spina-christi* mill. *J. Ethnopharmacol.* **52**:119–122.
32. Brinkworth, R. L., M. J. Stoermer, and D. P. Fairlie. 1992. Flavones are inhibitors of HIV-1 proteinase. *Biochem. Biophys. Res. Commun.* **188**:631–637.
33. Brownlee, H. E., A. R. McEuen, J. Hedger, and I. M. Scott. 1990. Antifungal effects of cocoa tannin on the witches' broom pathogen *Crinipellis perniciosus*. *Physiol. Mol. Plant Pathol.* **36**:39–48.
34. Burdick, E. M. 1971. Carpaine, an alkaloid of *Carica papaya*. Its chemistry and pharmacology. *Econ. Bot.* **25**:363–365.
35. Butler, L. G. 1988. Effects of condensed tannin on animal nutrition, p. 553. *In* R. W. Hemingway and J. J. Karchesy (ed.), *Chemistry and significance of condensed tannins*. Plenum Press, New York, N.Y.
36. Cardellina, J. H. I., M. H. G. Munro, R. W. Fuller, K. P. Manfredi, T. C. McKee, M. Tischler, H. R. Bokesch, K. R. Gustafson, J. A. Beutler, and M. R. Boyd. 1993. A chemical screening strategy for the dereplication and prioritization of HIV-inhibitory aqueous natural products extracts. *J. Nat. Prod.* **56**:1123–1129.
37. Casley-Smith, J. R., and J. R. Casley-Smith. 1997. Coumarin in the treatment of lymphoedema and other high-protein oedemas, p. 348. *In* R. O'Kennedy and R. D. Thornes (ed.), *Coumarins: biology, applications and mode of action*. John Wiley & Sons, Inc., New York, N.Y.
38. Chabot, S., R. Bel-Rhliid, R. Chenevert, and Y. Piche. 1992. Hyphal growth promotion in vitro of the VA mycorrhizal fungus, *Gigaspora margarita* Becker and Hall, by the activity of structurally specific flavonoid compounds under CO₂-enriched conditions. *New Phytol.* **122**:461–467.
39. Chaurasia, S. C., and K. K. Vyas. 1977. In vitro effect of some volatile oil against *Phytophthora parasitica* var. *piperina*. *J. Res. Indian Med. Yoga Homeopath.* **1977**:24–26.
40. Chen, K., Q. Shi, Y. Kashiwada, D. C. Zhang, C. Q. Hu, J. Q. Jin, H. Nozaki, R. E. Kilkuskie, E. Tramontano, Y. C. Cheng, et al. 1992. Anti-AIDS agents. 6. Salaspermic acid, an anti-HIV principle from *Tripterygium wilfordii*, and the structure-activity correlation with its related compounds. *J. Nat. Prod.* **55**:340–346.
41. Chessin, M., D. DeBorde, and A. Zipf (ed.). 1995. Antiviral proteins in higher plants. CRC Press, Inc., Boca Raton, Fla.
42. Cichewicz, R. H., and P. A. Thorpe. 1996. The antimicrobial properties of chile peppers (*Capsicum* species) and their uses in Mayan medicine. *J. Ethnopharmacol.* **52**:61–70.
43. Clark, A. M. 1996. Natural products as a resource for new drugs. *Pharm. Res.* **13**:1996.
44. Coe, S. D. 1994. America's first cuisines. University of Texas Press, Austin.
45. Colilla, F. J., A. Rocher, and E. Mendez. 1990. Gamma-purothionins: amino acid sequence of two polypeptides of a new family of thionins from wheat endosperm. *FEBS Lett.* **270**:191–194.
46. Corbeau, P., M. Haran, H. Binz, and C. Devaux. 1994. Jacalin, a lectin with anti-HIV-1 properties, and HIV-1 gp120 envelope protein interact with distinct regions of the CD4 molecule. *Mol. Immunol.* **31**:569–575.
47. Cordell, G. A., and O. E. Araujo. 1993. Capsaicin: identification, nomenclature, and pharmacotherapy. *Ann. Pharmacother.* **27**:330–336.
48. Critchfield, J. W., S. T. Butera, and T. M. Folks. 1996. Inhibition of HIV activation in latently infected cells by flavonoid compounds. *AIDS Res. Hum. Retroviruses* **12**:39.
49. Croom, E. M., Jr., and L. Walker. 1995. Botanicals in the pharmacy: new life for old remedies. *Drug Top.* **139**:84–93.
50. De Bolle, M. F., R. W. Osborn, I. J. Goderis, L. Noe, D. Acland, C. A. Hart, S. Torrekes, F. Van Leuven, and N. F. Broekart. 1996. Antimicrobial properties from *Mirabilis jalapa* and *Amaranthus caudatus*: expression, processing, localization and biological activity in transgenic tobacco. *Plant Mol. Biol.* **31**:993–1008.
51. De Clercq, E. 1995. Antiviral therapy for human immunodeficiency virus infections. *Clin. Microbiol. Rev.* **8**:200–239.
52. De Clercq, E. 1992. New perspectives for the chemotherapy of chemoprophylaxis of AIDS (acquired immunodeficiency syndrome). *Verhandelingen* **54**:57–89.
53. De Pasquale, R., M. P. Germano, A. Keita, R. Sanogo, and L. Iauk. 1995. Antiulcer activity of *Pteleopsis suberosa*. *J. Ethnopharmacol.* **47**:55–58.
54. Dhuley, J. 1998. Therapeutic efficacy of Ashwagandha against experimental aspergillosis in mice. *Immunopharmacol. Immunotoxicol.* **20**:191–198.
55. Di Stasi, L. C. 1995. Amoebicidal compounds from medicinal plants. *Parasitologia* **37**:29–39.
56. Dixon, R. A., P. M. Dey, and C. J. Lamb. 1983. Phytoalexins: enzymology and molecular biology. *Adv. Enzymol.* **55**:1–69.
57. Dornberger, K., V. Bockel, J. Heyer, C. Schonfeld, M. Tonew, and E. Tonew. 1975. Studies on the isothiocyanates erysolin and sulforaphan from *Cardaria draba*. *Pharmazie* **30**:792–796.
58. Duke, J. A. 1985. Handbook of medicinal herbs. CRC Press, Inc., Boca Raton, Fla.
59. Dwivedi, C., and A. Abu-Ghazaleh. 1997. Chemopreventive effects of sandalwood oil on skin papillomas in mice. *Eur. J. Cancer Prev.* **6**:399–401.
60. Eisenberg, D. M., R. C. Kessler, C. Foster, F. E. Norlock, D. R. Calkins, and T. L. Delbanco. 1993. Unconventional medicine in the United States: prevalence, costs and patterns of use. *N. Engl. J. Med.* **328**:246–252.
61. Eloff, J. N. 1998. Which extractant should be used for the screening and isolation of antimicrobial components from plants? *J. Ethnopharmacol.* **60**:1–8.
62. Estevez-Braun, A., R. Estevez-Reyes, L. M. Moujir, A. G. Ravelo, and A. G.

- Gonzalez.** 1994. Antibiotic activity and absolute configuration of 8S-hepta-deca-2(Z),9(Z)-diene-4,6-diyne-1,8-diol from *Bupleurum salicifolium*. *J. Nat. Prod.* **57**:1178–1182.
63. **Farley, D.** 1993. Making sure hype doesn't overwhelm science. *FDA Consumer* **27**:8–13.
64. **Favero, J., P. Corbeau, M. Nicolas, M. Benkirane, G. Trave, J. F. P. Dixon, P. Aucouturier, S. Rasheed, and J. W. Parker.** 1993. Inhibition of human immunodeficiency virus infection by the lectin jacalin and by a derived peptide showing a sequence similarity with GP120. *Eur. J. Immunol.* **23**:179–185.
65. **Fernandes de Caleyra, R., B. Gonzalez-Pascual, F. Garcia-Olmedo, and P. Carbonero.** 1972. Susceptibility of phytopathogenic bacteria to wheat purothionins in vitro. *Appl. Microbiol.* **23**:998–1000.
66. **Fernandez, M. A., M. D. Garcia, and M. T. Saenz.** 1996. Antibacterial activity of the phenolic acids fraction of *Scrophularia frutescens* and *Scrophularia sambucifolia*. *J. Ethnopharmacol.* **53**:11–14.
67. **Fesen, M. R., K. W. Kohn, F. Leteurtre, and Y. Pommier.** 1993. Inhibitors of human immunodeficiency virus integrase. *Proc. Natl. Acad. Sci. USA* **90**:2399–2403.
68. **Fesen, M. R., Y. Pommier, F. Leteurtre, S. Hiroguchi, J. Yung, and K. Kohn.** 1994. Inhibition of HIV-1 integrase by flavones, caffeic acid phenethyl ester (CAPE) and related compounds. *Biochem. Pharmacol.* **48**:595–608.
69. **Fessenden, R. J., and J. S. Fessenden.** 1982. Organic chemistry, 2nd ed. Willard Grant Press, Boston, Mass.
70. **Flayeh, K. A., and K. D. Sulayman.** 1987. Antimicrobial activity of the amine fraction of cucumber (*Cucumis sativus*) extract. *J. Appl. Microbiol.* **3**:275–279.
71. **Food and Drug Administration.** 1997. FDA talk paper: FDA publishes final dietary supplement rules, publication T97-45. Press Office, Food and Drug Administration, Washington, D.C.
72. **Francois, G., L. Ake Assi, J. Holenz, and G. Bringmann.** 1996. Constituents of *Picalima nitida* display pronounced inhibitory activities against asexual erythrocytic forms of *Plasmodium falciparum* in vitro. *J. Ethnopharmacol.* **54**:113–117.
73. **Freiburghaus, F., R. Kaminsky, M. H. H. Nkunya, and R. Brun.** 1996. Evaluation of African medicinal plants for their *in vitro* trypanocidal activity. *J. Ethnopharmacol.* **55**:1–11.
74. **Fujioka, T., and Y. Kashiwada.** 1994. Anti-AIDS agents. 11. Betulinic acid and platanic acid as anti-HIV principles from *Zygodium claviflorum*, and the anti-HIV activity of structurally related triterpenoids. *J. Nat. Prod.* **57**:243–247.
75. **Gautam, C. S., P. Pandhi, and P. L. Sharma.** 1993. Effect of oral administration of Abana, an ayurvedic formulation, on experimental cardiac arrhythmias. *Indian J. Pharmacol.* **2**:105–106.
76. **Geissman, T. A.** 1963. Flavonoid compounds, tannins, lignins and related compounds, p. 265. *In M. Florin and E. H. Stotz* (ed.), Pyrrole pigments, isoprenoid compounds and phenolic plant constituents, vol. 9. Elsevier, New York, N.Y.
77. **Georges, M., and K. M. Pandelai.** 1949. Investigations on plant antibiotics. IV. Further search for antibiotic substances in Indian medicinal plants. *Indian J. Med. Res.* **37**:169–181.
78. **Ghoshal, S., B. N. Krishna Prasad, and V. Lakshmi.** 1996. Antiamoebic activity of *Piper longum* fruits against *Entamoeba histolytica* in vitro and in vivo. *J. Ethnopharmacol.* **50**:167–170.
79. **Giron, L. M., G. A. Aguilar, A. Caceres, and G. L. Arroyo.** 1988. Anticandidal activity of plants used for the treatment of vaginitis in Guatemala and clinical trial of a *Solanum nigrescens* preparation. *J. Ethnopharmacol.* **22**:307–313.
80. **Gnabre, J. N., J. N. Brady, D. J. Clanton, Y. Ito, J. Dittmer, R. B. Bates, and R. C. C. Huang.** 1995. Inhibition of human immunodeficiency virus type 1 transcription and replication by DNA sequence-selective plant lignans. *Proc. Natl. Acad. Sci. USA* **92**:11239–11243.
81. **Gollapudi, S., H. A. Sharma, S. Aggarwal, L. D. Byers, H. E. Ensley, and S. Gupta.** 1995. Isolation of a previously unidentified polysaccharide (MAR-10) from *Hyssop officinalis* that exhibits strong activity against human immunodeficiency virus type 1. *Biochem. Biophys. Res. Commun.* **210**:145–151.
82. **Habtemariam, S., A. I. Gray, and P. G. Waterman.** 1993. A new antibacterial sesquiterpene from *Premna oligotricha*. *J. Nat. Prod.* **56**:140–143.
83. **Hamburger, H., and K. Hostettmann.** 1991. The link between phytochemistry and medicine. *Phytochemistry* **30**:3864–3874.
84. **Harrigan, G. G., A. Ahmad, N. Baj, T. E. Glass, A. A. L. Gunatilaka, and D. G. I. Kingston.** 1993. Bioactive and other sesquiterpenoids from *Porella cordeana*. *J. Nat. Prod.* **56**:921–925.
85. **Harris, R. S.** 1963. Vitamins K, p. 192–198. *In M. Florin and E. Stotz* (ed.), Pyrrole pigments, isoprenoid compounds and phenolic plant constituents, vol. 9. Elsevier, New York, N.Y.
86. **Hasegawa, H., S. Matsumiya, M. Uchiyama, T. Kurokawa, Y. Inouye, R. Kasai, S. Ishibashi, and K. Yamasaki.** 1994. Inhibitory effect of some triterpenoid saponins on glucose transport in tumor cells and its application to *in vitro* cytotoxic and antiviral activities. *Planta Med.* **6**:240–243.
87. **Haslam, E.** 1996. Natural polyphenols (vegetable tannins) as drugs: possible modes of action. *J. Nat. Prod.* **59**:205–215.
88. **Hayashi, K., M. Kamiya, and T. Hayashi.** 1995. Virucidal effects of the steam distillate from *Houttuynia cordata* and its components on HSV-1, influenza virus and HIV. *Planta Med.* **61**:237–241.
89. **Hess, S. C., R. L. Brum, N. K. Honda, A. B. Cruz, E. Moretto, R. B. Cruz, I. Messina, F. Ferrari, V. C. Filho, and R. A. Yunes.** 1995. Antibacterial activity and phytochemical analysis of *Vochysia divergens* (Vochysiaceae). *J. Ethnopharmacol.* **47**:97–100.
90. **Himejima, M., K. R. Hobson, T. Otsuka, D. L. Wood, and I. Kubo.** 1992. Antimicrobial terpenes from oleoresin of ponderosa pine tree *Pinus ponderosa*: a defense mechanism against microbial invasion. *J. Chem. Ecol.* **18**:1809–1818.
91. **Himejima, M., and I. Kubo.** 1991. Antibacterial agents from the cashew *Anacardium occidentale*. *J. Agric. Food Chem.* **39**:418–421.
92. **Holmes, O. W.** 1861. Currents and counter-currents in medical science, with other addresses and essays. Ticknor & Fields, Boston, Mass.
93. **Hopp, K. H., L. V. Cunningham, M. C. Bromel, L. J. Schermeister, and S. K. Wahba Khalil.** 1976. *In vitro* antitrypanosomal activity of certain alkaloids against *Trypanosoma lewisi*. *Lloydia* **39**:375–377.
94. **Houghton, P. J., T. Z. Woldemariam, A. I. Khan, A. Burke, and N. Mahmood.** 1994. Antiviral activity of natural and semi-synthetic chromosome alkaloids. *Antiviral Res.* **25**:235–244.
95. **Hoult, J. R. S., and M. Paya.** 1996. Pharmacological and biochemical actions of simple coumarins: natural products with therapeutic potential. *Gen. Pharmacol.* **27**:713–722.
96. **Hu, C. Q., K. Chen, Q. Shi, R. E. Kilkuskie, Y. C. Cheng, and K. H. Lee.** 1994. Anti-AIDS agents. 10. Acacetin-7-O-beta-D-galactopyranoside, and anti-HIV principle from *Chrysanthemum morifolium* and a structure-activity correlation with some related flavonoids. *J. Nat. Prod.* **57**:42–51.
97. **Huang, P. L., H. C. Chen, H. F. Kung, P. L. Huang, H. I. Huang, and S. Lee-Huang.** 1992. Anti-HIV plant proteins catalyze topological changes of DNA into inactive forms. *Biofactors* **4**:37–41.
98. **Hudson, J. B., F. Balza, L. Harris, and G. H. N. Towers.** 1993. Light-mediated activities of thiarubrines against human immunodeficiency virus. *Photochem. Photobiol.* **57**:675–680.
99. **Hudson, J. B., E. A. Graham, L. Harris, and M. J. Ashwood-Smith.** 1993. The unusual UVA-dependent antiviral properties of the furoisocoumarin, coufandrin. *Photochem. Photobiol.* **57**:491–496.
100. **Huffard, C. D., Y. Jia, E. M. Croom, Jr., I. Muhammed, A. L. Okunade, A. M. Clark, and R. D. Rogers.** 1993. Antimicrobial compounds from *Petalostemum purpureum*. *J. Nat. Prod.* **56**:1878–1889.
101. **Hunter, M. D., and L. A. Hull.** 1993. Variation in concentrations of phloridrin and phloretin in apple foliage. *Phytochemistry* **34**:1251–1254.
102. **Ito, M., H. Nakashima, M. Baba, R. Pauwels, E. De Clercq, S. Shigetani, and N. Yamamoto.** 1987. Inhibitory effect of glycyrrhizin on the *in vitro* infectivity and cytopathic activity of the human immunodeficiency virus [HIV (HTLV-III/LAV)]. *Antivir. Res.* **7**:127–137.
103. **Ivanovska, N., S. Philipov, R. Istatkova, and P. Georgieva.** 1996. Antimicrobial and immunological activity of ethanol extracts and fractions from *Isopyrum thalicroides*. *J. Ethnopharmacol.* **54**:143–151.
104. **Iwu, M. M., N. C. Unaeze, C. O. Okunji, D. G. Corley, D. R. Sanson, and M. S. Tempesta.** 1991. Antibacterial aromatic isothiocyanates from the essential oil of *Hippocratea welwitschii* roots. *Int. J. Pharmacogn.* **29**:154–158.
105. **Jones, G. A., T. A. McAllister, A. D. Muir, and K. J. Cheng.** 1994. Effects of sainfoin (*Onobrychis viciifolia* scop.) condensed tannins on growth and proteolysis by four strains of ruminal bacteria. *Appl. Environ. Microbiol.* **60**:1374–1378.
106. **Jones, N. L., S. Shabib, and P. M. Sherman.** 1997. Capsaicin as an inhibitor of the growth of the gastric pathogen *Helicobacter pylori*. *FEMS Microbiol. Lett.* **146**:223–227.
107. **Jones, S. B., Jr., and A. E. Luchsinger.** 1986. Plant systematics. McGraw-Hill Book Co., New York, N.Y.
108. **Kadota, S., P. Basnet, E. Ishii, T. Tamura, and T. Namba.** 1997. Antibacterial activity of trichorabdol from *Rabdosia trichocarpa* against *Helicobacter pylori*. *Zentbl. Bakteriol.* **286**:63–67.
109. **Kahane, I., and I. Ofek.** 1996. Toward anti-adhesion therapy for microbial diseases. Plenum Press, New York, N.Y.
110. **Kakiuchi, N., M. Hattori, and T. Namba.** 1985. Inhibitory effect of tannins on reverse transcriptase from RNA tumor virus. *J. Nat. Prod.* **48**:614–621.
111. **Kaul, T. N., E. Middletown, Jr., and P. L. Ogra.** 1985. Antiviral effect of flavonoids on human viruses. *J. Med. Virol.* **15**:71–79.
112. **Kazmi, M. H., A. Malik, S. Hameed, N. Akhtar, and S. Noor Ali.** 1994. An anthraquinone derivative from *Cassia italica*. *Phytochemistry* **36**:761–763.
113. **Keating, G. J., and R. O'Kennedy.** 1997. The chemistry and occurrence of coumarins, p. 348. *In R. O'Kennedy and R. D. Thornes* (ed.), Coumarins: biology, applications and mode of action. John Wiley & Sons, Inc., New York, N.Y.
114. **King, S. R., and M. S. Tempesta.** 1994. From shaman to human clinical trials: the role of industry in ethnobotany, conservation and community reciprocity. *Ciba Found. Symp.* **185**:197–206.
115. **Kitagawa, I., T. Mahmud, K.-I. Yokota, S. Nakagawa, T. Mayumi, M.**

- Kobayashi, and H. Shibuya. 1996. Indonesian medicinal plants. XVII. Characterization of quassinoids from the stems of *Quassia indica*. Chem. Pharm. Bull. **44**:2009–2014.
116. Klink, B. 1997. Alternative medicines: is natural really better? Drug Top. **141**:99–100.
117. Klopoukh, L., S. Siddiqi, M. Warns, and L. To. 1997. Growth inhibition of *Mycobacterium chelonae* and *Mycobacterium marinum* by plant compounds, abstr. A-199. In Abstracts of the 97th General Meeting of the American Society for Microbiology 1997. American Society for Microbiology, Washington, D.C.
118. Kragh, K. M., J. E. Nielsen, K. K. Nielsen, S. Dreboldt, and J. D. Mikkelsen. 1995. Characterization and localization of new antifungal cysteine-rich proteins from *Beta vulgaris*. Mol. Plant-Microbe Interact. **8**:424–434.
119. Kreis, W., M. H. Kaplan, J. Freeman, D. K. Sun, and P. S. Sarin. 1990. Inhibition of HIV replication by *Hyssop officinalis* extracts. Antiviral Res. **14**:323–337.
120. Kubo, A., C. S. Lunde, and I. Kubo. 1995. Antimicrobial activity of the olive oil flavor compounds. J. Agric. Food Chem. **43**:1629–1633.
121. Kubo, I., H. Muroi, and M. Himejima. 1992. Antibacterial activity of totarol and its potentiation. J. Nat. Prod. **55**:1436–1440.
122. Kubo, I., H. Muroi, and M. Himejima. 1993. Combination effects of antifungal naphthalenes against *Candida albicans* and two other fungi with phenylpropanoids. J. Nat. Prod. **56**:220–226.
123. Kubo, I., H. Muroi, and A. Kubo. 1994. Naturally occurring anti-acne agents. J. Nat. Prod. **57**:9–17.
124. Kuhnau, J. 1976. The flavonoids. A class of semi-essential food components: their role in human nutrition. World Rev. Nutr. Diet. **24**:117–191.
125. Kumar, O., and B. Singh. 1992. Effect of ayurvedic liver stimulants on live weight gain of broilers in north eastern region. Indian J. Anim. Res. **26**:1–5.
126. Lai, P. K., J. Donovan, H. Takayama, H. Sakagami, A. Tanaka, K. Konno, and M. Nonoyama. 1990. Modification of human immunodeficiency virus replication by pine cone extracts. AIDS Res. Hum. Retroviruses **6**:205–217.
127. Lee, C. D., M. Ott, S. P. Thyagarajan, D. A. Shafritz, R. D. Burk, and S. Gupta. 1996. *Phyllanthus amarus* down regulates hepatitis B virus mRNA transcription and replication. Eur. J. Clin. Invest. **26**:1069–1076.
128. Lee-Huang, S., P. L. Huang, H. C. Chen, P. L. Huang, A. Bourinbaiar, H. I. Huang, and H. F. Kung. 1995. Anti-HIV and anti-tumor activities of recombinant MAP30 from bitter melon. Gene (Amsterdam) **161**:151–156.
129. Lee-Huang, S., P. L. Huang, P. L. Huang, A. S. Bourinbaiar, H. C. Chen, and H. F. Kung. 1995. Inhibition of the integrase of human immunodeficiency virus (HIV) type 1 by anti-HIV plant proteins MAP30 and GAP31. Proc. Natl. Acad. Sci. USA **92**:8818–8822.
130. Lee-Huang, S., H. F. Kung, P. L. Huang, A. S. Bourinbaiar, J. L. Morell, J. H. Brown, P. L. Huang, W. P. Tsai, A. Y. Chen, H. I. Huang, and H. C. Chen. 1994. Human immunodeficiency virus type 1 (HIV-1) inhibition, DNA-binding, RNA-binding and ribosome inactivation activities in the N-terminal segments of the plant anti-HIV protein. Proc. Natl. Acad. Sci. USA **91**:12208–12212.
131. Lewis, W. H., and M. P. Elvin-Lewis. 1995. Medicinal plants as sources of new therapeutics. Ann. Mo. Bot. Gard. **82**:16–24.
132. Liener, I. E. 1970. Toxic constituents of plant foodstuffs. Proc. Nutr. Soc. **29**:56–57.
133. Manonmani, S., V. P. Vishwanathan, S. Subramanian, and S. Govindasamy. 1995. Biochemical studies on the antitumor activity of Cauvery 100, an ayurvedic formulation in experimental ulcers. Indian J. Pharmacol. **27**:101–105.
134. Manonmani, S., S. William, S. Subramanian, and S. Govindasamy. 1991. Biochemical studies on the antidiarrhoeal effects of Cauvery-100, an ayurvedic formulation, in rats. Biochem. Int. **24**:701–708.
135. Martin, G. J. 1995. Ethnobotany: a methods manual. Chapman & Hall, New York, N.Y.
136. Martinez, M. J., J. Betancourt, N. Alonso-Gonzalez, and A. Jauregui. 1996. Screening of some Cuban medicinal plants for antimicrobial activity. J. Ethnopharmacol. **52**:171–174.
137. Mason, T. L., and B. P. Wasserman. 1987. Inactivation of red beet beta-glucan synthase by native and oxidized phenolic compounds. Phytochemistry **26**:2197–2202.
138. Mayr-Harting, A., A. Hedges, and R. Berkelery (ed.). 1972. Methods for studying bacteriocins. Academic Press, Inc., New York, N.Y.
139. Mazumder, A., D. Cooney, R. Agbaria, and Y. Pommier. 1994. Inhibition of human immunodeficiency virus type 1 integrase by 3'-azido-3'-deoxythymidylate. Proc. Natl. Acad. Sci. USA **91**:5771–5775.
140. McDevitt, J. T., D. M. Schneider, S. K. Katiyar, and T. D. Edlind. 1996. Berberine: a candidate for the treatment of diarrhea in AIDS patients, abstr. 175. In Program and Abstracts of the 36th Interscience Conference on Antimicrobial Agents and Chemotherapy. American Society for Microbiology, Washington, D.C.
141. McGrath, M. S., K. M. Hwang, S. E. Caldwell, I. Gaston, K. C. Luk, P. Wu, V. L. Ng, S. Crowe, J. Daniels, J. Marsh, et al. 1989. GLQ223: an inhibitor of human immunodeficiency virus replication in acutely and chronically infected cells of lymphocyte and mononuclear phagocyte lineage. Proc. Natl. Acad. Sci. USA **86**:2844–2848.
142. McMahon, J. B., M. J. Currens, R. J. Gulakowski, R. W. J. Buckheit, C. Lackman-Smith, Y. F. Hallock, and M. R. Boyd. 1995. Michellamine B, a novel plant alkaloid, inhibits human immunodeficiency virus-induced cell killing by at least two distinct mechanisms. Antimicrob. Agents Chemother. **39**:484–488.
143. Mendez, E., A. Moreno, F. Colilla, F. Pelaez, G. G. Limas, R. Mendez, F. Soriano, M. Salinas, and C. De Haro. 1990. Primary structure and inhibition of protein synthesis in eukaryotic cell-free system of a novel thionin, gamma-horothionin, from barley endosperm. Eur. J. Biochem. **194**:533–539.
144. Mendoza, L., M. Wilkens, and A. Urzua. 1997. Antimicrobial study of the resinous exudates and of diterpenoids and flavonoids isolated from some Chilean *Pseudognaphalium* (Asteraceae). J. Ethnopharmacol. **58**:85–88.
145. Meyer, J. J. M., A. J. Afolayan, M. B. Taylor, and D. Erasmus. 1997. Antiviral activity of galangin from the aerial parts of *Helichrysum aureonitens*. J. Ethnopharmacol. **56**:165–169.
146. Moerman, D. E. 1996. An analysis of the food plants and drug plants of native North America. J. Ethnopharmacol. **52**:1–22.
147. Moore, P. S., and C. Pizza. 1992. Observations on the inhibition of HIV-1 reverse transcriptase by catechins. Biochem. J. **288**:717–719.
148. Muller, W. E. G., K. Renneisen, M. H. Kreuter, H. C. Schroder, and I. Winkler. 1988. The D-mannose-specific lectin from *Gerardia savaglia* blocks binding of human immunodeficiency virus type I to H9 cells and human lymphocytes in vitro. J. Acquired Immune Defic. Syndr. **1**:453–458.
149. Murakami, A., H. Ohgashi, S. Tanaka, A. Tatematsu, and K. Koshimizu. 1993. Bitter cyanoglucosides from *Lophira alata*. Phytochemistry **32**:1461–1466.
150. Naganawa, R., N. Iwata, K. Ishikawa, H. Fukuda, T. Fujino, and A. Suzuki. 1996. Inhibition of microbial growth by ajoene, a sulfur-containing compound derived from garlic. Appl. Environ. Microbiol. **62**:4238–4242.
151. Nakahara, K., S. Kawabata, H. Ono, K. Ogura, T. Tanaka, T. Ooshima, and S. Hamada. 1993. Inhibitory effect of oolong tea polyphenols on glucosyltransferases of mutans streptococci. Appl. Environ. Microbiol. **59**:968–973.
152. Namba, T., O. Morita, S.-L. Huang, K. Goshima, M. Hattori, and N. Kakiuchi. 1988. Studies on cardio-active crude drugs. I. Effect of coumarins on cultured myocardial cells. Planta Med. **54**:277–282.
153. Navarro, V., M. L. Villarreal, G. Rojas, and X. Lozoya. 1996. Antimicrobial evaluation of some plants used in Mexican traditional medicine for the treatment of infectious diseases. J. Ethnopharmacol. **53**:143–147.
154. Ngan, F., R. S. Chang, H. D. Tabba, and K. M. Smith. 1988. Isolation, purification and partial characterization of an active anti-HIV compound from the Chinese medicinal herb *Viola yedoensis*. Antiviral Res. **10**:107–116.
155. Nonaka, G.-I., I. Nishioka, M. Nishizawa, T. Yamagishi, Y. Kashiwada, G. E. Dutschman, A. J. Bodner, R. E. Killuskie, Y.-C. Cheng, and K.-H. Lee. 1990. Anti-AIDS agents. 2. Inhibitory effects of tannins on HIV reverse transcriptase and HIV replication in H9 lymphocyte cells. J. Nat. Prod. **53**:587–595.
156. Norton, M. R., and M. Addy. 1989. Chewing sticks versus toothbrushes in West Africa. A pilot study. Clin. Prev. Dent. **11**:311–313.
157. Odebiyi, O. O., and E. A. Sofowora. 1979. Antimicrobial alkaloids from a Nigerian chewing stick (*Fagara zanthoxyloides*). Planta Med. **36**:204–207.
158. Ofek, I., J. Goldhar, and N. Sharon. 1996. Anti-*Escherichia coli* adhesion activity of cranberry and blueberry juices. Adv. Exp. Med. Biol. **408**:179–183.
159. Ofek, I., J. Goldhar, D. Zafiriri, H. Lis, R. Adar, and N. Sharon. 1991. Anti-*Escherichia coli* adhesion activity of cranberry and blueberry juices. N. Engl. J. Med. **324**:1599. (Letter.)
160. Offergeld, R., C. Reinecker, E. Gumz, S. Schrum, R. Treiber, R. D. Neth, and S. H. Gohla. 1992. Mitogenic activity of high molecular polysaccharide fractions isolated from the cupressaceae *Thuja occidentalis* L. enhanced cytokine-production by thypolysaccharide, g-fraction (TPSG). Leukemia **6**(Suppl. 3):189S–191S.
161. O'Kennedy, R., and R. D. Thornes (ed.). 1997. Coumarins: biology, applications and mode of action. John Wiley & Sons, Inc., New York, N.Y.
162. Oliver-Bever, B. 1986. Medicinal plants in tropical West Africa. Cambridge University Press, New York, N.Y.
163. Omulokoli, E., B. Khan, and S. C. Chhabra. 1997. Antiplasmodial activity of four Kenyan medicinal plants. J. Ethnopharmacol. **56**:133–137.
164. Ono, K., H. Nakane, M. Fukushima, J.-C. Chermann, and F. Barre-Sinoussi. 1989. Inhibition of reverse transcriptase activity by a flavonoid compound, 5,6,7-trihydroxyflavone. Biochem. Biophys. Res. Commun. **3**:982–987.
165. Ono, K., H. Nakane, M. Fukushima, J.-C. Chermann, and F. Barre-Sinoussi. 1990. Differential inhibitory effects of various flavonoids on the activities of reverse transcriptase and cellular DNA and RNA polymerases. Eur. J. Biochem. **190**:469–476.
166. Ooshima, T., T. Minami, W. Aono, A. Izumitani, S. Sobue, T. Fujiwara, S. Kawabata, and S. Hamada. 1993. Oolong tea polyphenols inhibit experimental dental caries in SPF rats infected with mutans streptococci. Caries Res. **27**:124–129.

167. Orozco-Topete, R., J. Sierra-Madero, C. Cano-Dominguez, J. Kershenvich, G. Ortiz-Pedroza, E. Vazquez-Valls, C. Garcia-Cosio, A. Soria-Cordoba, A. M. Armendariz, X. Teran-Toledo, J. Romo-Garcia, H. Fernandez, and E. J. Rozhon. 1997. Safety and efficacy of Virend for topical treatment of genital and anal herpes simplex lesions in patients with AIDS. *Antiviral Res.* 35:91-103.
168. Osato, J. A., L. A. Santiago, G. M. Remo, M. S. Cuadra, and A. Mori. 1993. Antimicrobial and antioxidant activities of unripe papaya. *Life Sci.* 53:1383-1389.
169. Pantazis, P. 1996. Camptothecin: a promising antiretroviral drug. *J. Biomed. Sci.* 3:14-19.
170. Paranjpe, P., and P. H. Kulkarni. 1995. Comparative efficacy of four Ayurvedic formulations in the treatment of acne vulgaris: a double-blind randomised placebo-controlled clinical evaluation. *J. Ethnopharmacol.* 49:127-132.
171. Paris, A., B. Strukelj, M. Renko, and V. Turk. 1993. Inhibitory effect of carnosolic acid on HIV-1 protease in cell-free assays. *J. Nat. Prod.* 56:1426-1430.
172. Pengsuparp, T., L. Cai, H. Constant, H. H. Fong, L. Z. Lin, A. D. Kinghorn, J. M. Pezzuto, G. A. Cordell, K. Ingoldsdottir, and H. Wagner. 1995. Mechanistic evaluation of new plant-derived compounds that inhibit HIV-1 reverse transcriptase. *J. Nat. Prod.* 58:1024-1031.
173. Pengsuparp, T., L. Cai, H. H. S. Fong, A. D. Kinghorn, J. M. Pezzuto, M. C. Wani, and M. E. Wall. 1994. Pentacyclic triterpenes derived from *Maprounea africana* are potent inhibitors of HIV-1 reverse transcriptase. *J. Nat. Prod.* 57:415-418.
174. Peres, M. T. L. P., F. D. Monache, A. B. Cruz, M. G. Pizzolatti, and R. A. Yunes. 1997. Chemical composition and antimicrobial activity of *Croton urucurana* Baillon (Euphorbiaceae). *J. Ethnopharmacol.* 56:223-226.
175. Perrett, S., P. J. Whitfield, L. Sanderson, and A. Bartlett. 1995. The plant molluscicide *Millettia thonningii* (Leguminosae) as a topical antischistosomal agent. *J. Ethnopharmacol.* 47:49-54.
176. Phillipson, J. D., and M. J. O'Neill. 1987. New leads to the treatment of protozoal infections based on natural product molecules. *Acta Pharm. Nord.* 1:131-144.
177. Piller, N. B. 1975. A comparison of the effectiveness of some anti-inflammatory drugs on thermal oedema. *Br. J. Exp. Pathol.* 56:554-559.
178. Prpic-Majic, D., A. Pizent, J. Jurasovic, J. Pongracic, and N. Restek-Samarzija. 1996. Lead poisoning associated with the use of Ayurvedic metal-mineral tonics. *J. Toxicol. Clin. Toxicol.* 34:417-423.
179. Rana, B. K., U. P. Singh, and V. Taneja. 1997. Antifungal activity and kinetics of inhibition by essential oil isolated from leaves of *Aegle marmelos*. *J. Ethnopharmacol.* 57:29-34.
180. Rao, K. V., K. Sreeramulu, D. Gunasekar, and D. Ramesh. 1993. Two new sesquiterpene lactones from *Ceiba pentandra*. *J. Nat. Prod.* 56:2041-2045.
181. Rinehart, K. L., T. G. Holt, N. L. Fregeau, P. A. Keifer, G. R. Wilson, T. J. Perum, Jr., R. Sakai, A. G. Thompson, J. G. Stroh, L. S. Shield, and D. S. Seigler. 1990. Bioactive compounds from aquatic and terrestrial sources. *J. Nat. Prod.* 53:771-792.
182. Rojas, A., L. Hernandez, R. Pereda-Miranda, and R. Mata. 1992. Screening for antimicrobial activity of crude drug extracts and pure natural products from Mexican medicinal plants. *J. Ethnopharmacol.* 35:275-283.
183. Rotimi, V. O., B. E. Laughon, J. G. Bartlett, and H. A. Mosadami. 1988. Activities of Nigerian chewing stick extracts against *Bacteroides gingivalis* and *Bacteroides melaninogenicus*. *Antimicrob. Agents Chemother.* 32:598-600.
184. Rucker, G., S. Kehrbaum, H. Sakulas, B. Lawong, and F. Goeltenboth. 1992. Acetylenic glucosides from *Microglossa pyrifolia*. *Planta Med.* 58:266-269.
185. Sakanaka, S., M. Kim, M. Taniguchi, and T. Yamamoto. 1989. Antibacterial substances in Japanese green tea extract against *Streptococcus mutans*, a cariogenic bacterium. *Agric. Biol. Chem.* 53:2307-2311.
186. Sakanaka, S., N. Shimura, M. Aizawa, M. Kim, and T. Yamamoto. 1992. Preventive effect of green tea polyphenols against dental caries in conventional rats. *Biosci. Biotechnol. Biochem.* 56:592-594.
187. San-Blas, G., L. Marino, F. San-Blas, and R. Apitz-Castro. 1993. Effect of ajoene on dimorphism of *Paracoccidioides brasiliensis*. *J. Med. Vet. Mycol.* 31:133-141.
188. San-Blas, G., F. San-Blas, F. Gil, L. Marino, and R. Apitz-Castro. 1989. Inhibition of growth of the dimorphic fungus *Paracoccidioides brasiliensis* by ajoene. *Antimicrob. Agents Chemother.* 33:1641-1644.
189. Sato, M., S. Fujiwara, H. Tsuchiya, T. Fujii, M. Inuma, H. Tosa, and Y. Ohkawa. 1996. Flavones with antibacterial activity against cariogenic bacteria. *J. Ethnopharmacol.* 54:171-176.
190. Sato, Y., H. Odetani, K. Singyouchi, T. Ohtsubo, M. Kihara, H. Shibata, and T. Higuti. 1997. Extraction and purification of effective antimicrobial constituents of *Terminalia chebula* RETS against methicillin-resistant *Staphylococcus aureus*. *Biol. Pharm. Bull.* 20:401-404.
191. Satrija, F., P. Nansen, S. Murtini, and S. He. 1995. Anthelmintic activity of papaya latex against patent *Heligmosomoides polygyrus* infections in mice. *J. Ethnopharmacol.* 48:161-164.
192. Scalbert, A. 1991. Antimicrobial properties of tannins. *Phytochemistry* 30:3875-3883.
193. Scheel, L. D. 1972. The biological action of the coumarins. *Microbiol. Toxins* 8:47-66.
194. Schmidt, H. 1988. Phenol oxidase (E.I.14.18.1), a marker enzyme for defense cells. Progress in histochemistry and cytochemistry, vol. 17. Gustav Fischer, New York, N.Y.
195. Schultes, R. E. 1978. The kingdom of plants, p. 208. In W. A. R. Thomson (ed.), *Medicines from the Earth*. McGraw-Hill Book Co., New York, N.Y.
196. Schultz, J. C. 1988. Tannin-insect interactions, p. 553. In R. W. Hemingway and J. J. Karchesy (ed.), *Chemistry and significance of condensed tannins*. Plenum Press, New York, N.Y.
197. Scortichini, M., and M. Pia Rossi. 1991. Preliminary *in vitro* evaluation of the antimicrobial activity of terpenes and terpenoids towards *Erwinia amylovora* (Burrill) Winslow *et al.* *J. Appl. Bacteriol.* 71:109-112.
198. Serafini, M., A. Ghiselli, and A. Ferro-Luzzi. 1994. Red wine, tea and anti-oxidants. *Lancet* 344:626.
199. Serkedjjeva, J., and N. Manolova. 1992. Anti-influenza virus effect of some propolis constituents and their analogues (esters of substituted cinnamic acids). *J. Nat. Prod.* 55:294-297.
200. Sethi, M. L. 1979. Inhibition of reverse transcriptase activity by benzophenanthridine alkaloids. *J. Nat. Prod.* 42:187-196.
201. Shah, L., S. P. Patil, and J. Patil. 1997. Observations on clinical evaluation of indigenous herbal drugs in the treatment of mental illnesses. *Indian J. Pharmacol.* 29:S347-S349.
202. Sharon, N., and I. Ofek. 1986. Mannose specific bacterial surface lectins, p. 55-82. In D. Mirelman (ed.), *Microbial lectins and agglutinins*. John Wiley & Sons, Inc., New York, N.Y.
203. Silva, O., A. Duarte, J. Cabrita, M. Pimentel, A. Diniz, and E. Gomes. 1996. Antimicrobial activity of Guinea-Bissau traditional remedies. *J. Ethnopharmacol.* 50:55-59.
204. Silva, O., A. Duarte, M. Pimental, S. Viegas, H. Barroso, J. Machado, I. Pires, J. Cabrita, and E. Gomes. 1997. Antimicrobial activity of *Terminalia macroptera* root. *J. Ethnopharmacol.* 57:203-207.
205. Sohni, Y. R., P. Kaimal, and R. M. Bhatt. 1995. The antiamebic effect of a crude drug formation of herbal extracts against *Entamoeba histolytica* in vitro and in vivo. *J. Ethnopharmacol.* 45:43-52.
206. Soine, T. O. 1964. Naturally occurring coumarins and related physiological activities. *J. Pharm. Sci.* 53:231-264.
207. Song, Z. J., H. K. Johansen, V. Faber, and N. Hoiby. 1997. Ginseng treatment enhances bacterial clearance and decreases lung pathology in athymic rats with chronic *Pseudomonas aeruginosa* pneumonia. *APMIS* 105:438-444.
208. Spedding, G., A. Ratty, and E. J. Middleton. 1989. Inhibition of reverse transcriptases by flavonoids. *Antiviral Res.* 12:99-110.
209. Stange, R. R., Jr., S. L. Midland, J. W. Eckert, and J. J. Sims. 1993. An antifungal compound produced by grapefruit and Valencia orange after wounding of the peel. *J. Nat. Prod.* 56:1627-1629.
210. Stern, J. L., A. E. Hagerman, P. D. Steinberg, and P. K. Mason. 1996. Phlorotannin-protein interactions. *J. Chem. Ecol.* 22:1887-1899.
211. Stockwell, C. 1988. *Nature's pharmacy*. Century Hutchinson Ltd., London, United Kingdom.
212. Sun, H. D., S. X. Qiu, L. Z. Lin, Z. Y. Wang, Z. W. Lin, T. Pengsuparp, J. M. Pezzuto, H. H. Fong, G. A. Cordell, and N. R. Farnsworth. 1996. Nigranoic acid, a triterpenoid from *Schisandra sphaerandra* that inhibits HIV-1 reverse transcriptase. *J. Nat. Prod.* 59:525-527.
213. Suresh, B., S. Sriram, S. A. Dhanaraj, K. Elango, and K. Chinnaaswamy. 1997. Anticandidal activity of *Santolina chamaecyparissus* volatile oil. *J. Ethnopharmacol.* 55:151-159.
214. Tabba, H. D., R. S. Chang, and K. M. Smith. 1989. Isolation, purification, and partial characterization of prunellin, an anti-HIV component from aqueous extracts of *Prunella vulgaris*. *Antiviral Res.* 11:263-273.
215. Tada, M., Y. Hiroe, S. Kiyohara, and S. Suzuki. 1988. Nematicidal and antimicrobial constituents from *Allium grayi* Regel and *Allium fistulosum* L. var. *caespitosum*. *Agric. Biol. Chem.* 52:2383-2385.
216. Tan, G. T., J. F. Miller, A. D. Kinghorn, S. H. Hughes, and J. M. Pezzuto. 1992. HIV-1 and HIV-2 reverse transcriptases: a comparative study of sensitivity to inhibition by selected natural products. *Biochem. Biophys. Res. Commun.* 185:370-378.
217. Tan, G. T., J. M. Pezzuto, and A. D. Kinghorn. 1991. Evaluation of natural products as inhibitors of human immunodeficiency virus type 1 (HIV-1) reverse transcriptase. *J. Nat. Prod.* 54:143-154.
218. Tang, R., K. Chen, M. Cosentino, and K.-H. Lee. 1994. Apigenin-7-O- β -D-glucopyranoside, an anti-HIV principle from *Kummerowia striata*. *Bioorg. Med. Chem. Lett.* 4:455-458.
219. Taniguchi, M., and I. Kubo. 1993. Ethnobotanical drug discovery based on medicine men's trials in the African savanna: screening of east African plants for antimicrobial activity II. *J. Nat. Prod.* 56:1539-1546.
220. Tassou, C. C., E. H. Drosinos, and G. J. E. Nychas. 1995. Effects of essential oil from mint (*Mentha piperita*) on *Salmonella enteritidis* and *Listeria monocytogenes* in model food systems at 4° and 10°C. *J. Appl. Bacteriol.* 78:593-600.

221. Taylor, R. S. L., F. Edel, N. P. Manandhar, and G. H. N. Towers. 1996. Antimicrobial activities of southern Nepalese medicinal plants. *J. Ethnopharmacol.* **50**:97–102.
222. Terras, F. R. G., H. M. E. Schoofs, H. M. E. Thevissen, R. W. Osborn, J. Vanderleyden, B. P. A. Cammue, and W. F. Broekaert. 1993. Synergistic enhancement of the antifungal activity of wheat and barley thionins by radish and oilseed rape 2S albumins and by barley trypsin inhibitors. *Plant Physiol.* **103**:1311–1319.
223. Thastrup, O., J. B. Knudsen, J. Lemmich, and K. Winther. 1985. Inhibitions of human platelet aggregation by dihydropyran- and dihydrofuranocoumarins, a new class of cAMP phosphodiesterase inhibitors. *Biochem. Pharmacol.* **34**:2137–2140.
224. Thomson, W. A. R. (ed.). 1978. *Medicines from the Earth*. McGraw-Hill Book Co., Maidenhead, United Kingdom.
225. Thornes, R. D. 1997. Clinical and biological observations associated with coumarins, p. 256. In R. O'Kennedy and R. D. Thornes (ed.), *Coumarins: biology, applications and mode of action*. John Wiley & Sons, Inc., New York, N.Y.
226. Toda, M., S. Okubo, H. Ikigai, T. Suzuki, Y. Suzuki, Y. Hara, and T. Shimamura. 1992. The protective activity of tea catechins against experimental infection by *Vibrio cholerae* O1. *Microbiol. Immunol.* **36**:999–1001.
227. Toda, M., S. Okubo, R. Ohnishi, and T. Shimamura. 1989. Antibacterial and bactericidal activities of Japanese green tea. *Jpn. J. Bacteriol.* **45**:561–566.
228. Tsuchiya, H., M. Sato, M. Iinuma, J. Yokoyama, M. Ohyama, T. Tanaka, I. Takase, and I. Namikawa. 1994. Inhibition of the growth of cariogenic bacteria in vitro by plant flavanones. *Experientia* **50**:846–849.
229. Tsuchiya, H., M. Sato, T. Miyazaki, S. Fujiwara, S. Tanigaki, M. Ohyama, T. Tanaka, and M. Iinuma. 1996. Comparative study on the antibacterial activity of phytochemical flavanones against methicillin-resistant *Staphylococcus aureus*. *J. Ethnopharmacol.* **50**:27–34.
230. Tyler, V. E. 1987. *The new honest herbal*. George F. Stickley Co., Philadelphia, Pa.
231. Urs, N. V. R. R., and J. M. Dunleavy. 1975. Enhancement of the bactericidal activity of a peroxidase system by phenolic compounds (*Xanthomonas phaseoli* var. *sojensis*, soybeans). *Phytopathology* **65**:686–690.
232. U.S. Department of Health and Human Services. 1992. National Toxicology Program technical report on the toxicology and carcinogenesis of coumarin in F344/N rats and B6C3F1 mice (gavage studies). NIH publication 92-3153. U.S. Department of Health and Human Services, Washington, D.C.
233. Vamos-Vigyazo, L. 1981. Polyphenol oxidase and peroxidase in fruits and vegetables. *Crit. Rev. Food Sci. Nutr.* **15**:49–127.
234. Vanden Berghe, D. A., A. J. Vlietinck, and L. Van Hoof. 1986. Plant products as potential antiviral agents. *Bull. Inst. Pasteur* **84**:101–147.
235. Vijaya, K., S. Ananthan, and R. Nalini. 1995. Antibacterial effect of theaflavin, polyphenon 60 (*Camellia sinensis*) and *Euphorbia hirta* on *Shigella* spp.—a cell culture study. *J. Ethnopharmacol.* **49**:115–118.
236. Virus, R. M., and G. F. Gebhart. 1979. Pharmacologic actions of capsaicin: apparent involvement of substance P and serotonin. *Life Sci.* **25**:1273–1284.
237. Vishwakarma, R. A. 1990. Stereoselective synthesis of α -arteether from artemisinin. *J. Nat. Prod.* **53**:216–217.
238. Vlietinck, A. J., and D. A. Vanden Berghe. 1991. Can ethnopharmacology contribute to the development of antiviral drugs? *J. Ethnopharmacol.* **32**:141–153.
239. Vohora, S. B., M. Rizwan, and J. A. Khan. 1973. Medicinal uses of common Indian vegetables. *Planta Med.* **23**:381–393.
240. Walker, C. B. 1988. Microbiological effects of mouthrinses containing antimicrobials. *J. Clin. Periodontol.* **15**:499–505.
241. Wan, J., A. Wilcock, and M. J. Coventry. 1998. The effect of essential oils of basil on the growth of *Aeromonas hydrophila* and *Pseudomonas fluorescens*. *J. Appl. Microbiol.* **84**:152–158.
242. Watanbe, H., C. Miyaji, M. Makino, and T. Abo. 1996. Therapeutic effects of glycyrrhizin in mice infected with LP-BM5 murine retrovirus and mechanisms involved in the prevention of disease progression. *Biotherapy* **9**:209–220.
243. Weiner, M. A. 1980. *Earth medicine—earth food: plant remedies, drugs and natural foods of the North American Indians*. Macmillan, New York, N.Y.
244. Weimann, I. 1997. History of the development and applications of coumarin and coumarin-related compounds. In R. O'Kennedy and R. D. Thornes (ed.), *Coumarins: biology, applications and mode of action*. John Wiley & Sons, Inc., New York, N.Y.
245. Wild, R. (ed.). 1994. *The complete book of natural and medicinal cures*. Rodale Press, Inc., Emmaus, Pa.
246. Xu, H. X., F. Q. Zeng, M. Wan, and K. Y. Sim. 1996. Anti-HIV triterpene acids from *Geum japonicum*. *J. Nat. Prod.* **59**:643–645.
247. Ya, C., S. H. Gaffney, T. H. Lilley, and E. Haslam. 1988. Carbohydrate-polyphenol complexation, p. 553. In R. W. Hemingway and J. J. Karchesy (ed.), *Chemistry and significance of condensed tannins*. Plenum Press, New York, N.Y.
248. Yankauer, A. 1997. The recurring popularity of alternative medicine. *Perspect. Biol. Med.* **41**:132–137.
249. Yao, X.-J., M. A. Wainberg, and M. A. Parniak. 1992. Mechanism of inhibition of HIV-1 infection in vitro by purified extract of *Prunella vulgaris*. *Virology* **178**:56.
250. Yoshida, S., S. Kasuga, N. Hayashi, T. Ushiroguchi, H. Matsuura, and S. Nakagawa. 1987. Antifungal activity of ajoene derived from garlic. *Appl. Environ. Microbiol.* **53**:615–617.
251. Yoshikawa, M., E. Harada, Y. Naitoh, K. Inoue, H. Matsuda, H. Shimoda, J. Yamahara, and N. Murakami. 1994. Development of bioactive functions in *Hydrangeae dulcis* folium. III. On the anti-allergic and antimicrobial principles of *Hydrangeae dulcis* folium. (1) Thunberginsols A, B and F. *Chem. Pharm. Bull. (Tokyo)* **42**:2225–2230.
252. Zafri, D., I. Ofek, R. Adar, M. Pocino, and N. Sharon. 1989. Inhibitory activity of cranberry juice on adherence of type 1 and type P fimbriated *Escherichia coli* to eucaryotic cells. *Antimicrob. Agents Chemother.* **33**:92–98.
253. Zhang, Y., and K. Lewis. 1997. Fabatins: new antimicrobial plant peptides. *FEMS Microbiol. Lett.* **149**:59–64.